

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 December 2001 (20.12.2001)

PCT

(10) International Publication Number
WO 01/96584 A2

(51) International Patent Classification⁷: **C12N 15/82**

(21) International Application Number: PCT/US01/18911

(22) International Filing Date: 12 June 2001 (12.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/210,917 12 June 2000 (12.06.2000) US

(71) Applicant (for all designated States except US):
AKKADIX CORPORATION [US/US]; 4204 Sorrento Valley Blvd., Suite A, San Diego, CA 92121-1412 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MUSHEGIAN, Arcady, R.** [—/US]; 3987 Santa Nella Place, San Diego, CA 92130 (US). **TAYLOR, Christopher, G.** [—/US]; 2910-A

Luciemaga Street, Carlsbad, CA 92009 (US). **FEITELSON, Jerald, S.** [—/US]; 4387 Mistral Place, San Diego, CA 92130 (US). **EROSHKIN, Alexy, M.** [—/US]; 3803 Ruette San Rafael, San Diego, CA 92130 (US).


(74) Agents: **LLOYD, Jeff** et al.; Saliwanchik, Lloyd & Saliwanchik, Suite A-1, 2421 N.W. 41st Street, Gainesville, FL 32606 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES



(57) Abstract: The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides RNAi molecules, polynucleotide sequences, and methods of using these sequences in nematode control.

WO 01/96584 A2



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

DESCRIPTION

MATERIALS AND METHODS FOR THE CONTROL OF NEMATODES

Background of the Invention

[0001] Plant parasitic nematodes, such as root-knot nematodes (*Meloidogyne* species) and cyst nematodes (*Globodera* and *Heterodera*), attack nearly every food crop, and are among the world's most damaging agricultural pests. For example, root-knot nematodes parasitize more than 2,000 plant species from diverse plant families and represent a tremendous threat to crop production world-wide. These biotrophic pathogens have evolved highly specialized and complex feeding relationships with their hosts.

[0002] Nematodes cause millions of dollars of damage each year to turf grasses, ornamental plants, and food crops. Efforts to eliminate or minimize damage caused by nematodes in agricultural settings have typically involved the use of soil fumigation with materials such as chloropicrin, methyl bromide, and dazomet, which volatilize to spread the active ingredient throughout the soil. Such fumigation materials can be highly toxic and may create an environmental hazard. Various non-fumigant chemicals have also been used, but these too create serious environmental problems and can be highly toxic to humans.

[0003] Some research articles have been published concerning the effects of δ -endotoxins from *B. thuringiensis* species on the viability of nematodes. See, for example, Bottjer, Bone and Gill ([1985] *Experimental Parasitology* 60:239-244); Ignoffo and Dropkin (Ignoffo, C.M., Dropkin, V.H. [1977] *J. Kans. Entomol. Soc.* 50:394-398); and Ciordia, H. and W.E. Bizzell ([1961] *Jour. of Parasitology* 47:41 [abstract]). Several patents have issued describing the control of nematodes with *B.t.* See, for example, U.S. Patent Nos. 4,948,734; 5,093,120; 5,281,530; 5,426,049; 5,439,881; 5,236,843; 5,322,932; 5,151,363; 5,270,448; 5,350,577; 5,667,993; and 5,670,365. The development of resistance by insects to *B.t.* toxins is one obstacle to the successful use of such toxins.

[0004] The pesticidal activity of avermectins is well known. The avermectins are disaccharide derivatives of pentacyclic, 16-membered lactones. They can be divided into four major compounds: A_{1a}, A_{2a}, B_{1a}, and B_{2a}; and four minor compounds: A_{1b}, A_{2b}, B_{1b}, and B_{2b}. The isolation and purification of these compounds is also described in U.S. Patent No. 4,310,519, issued January 12, 1982. Avermectin B_{2a} is active against the root-knot nematode, *Meloidogyne incognita*. It is reported to be 10-30 times as potent as commercial contact nematicides when incorporated into soil at 0.16-0.25 kg/ha (Boyce Thompson Institute for Plant Research 58th Annual Report [1981]; Putter, I. *et al.* [1981] "Avermectins: Novel Insecticides, Acaracides, and Nematicides from a Soil Microorganism," *Experientia* 37:963-964). Avermectin B_{2a} is not toxic to tomatoes or cucumbers at rates of up to 10 kg/ha.

[0005] Fatty acids are a class of natural compounds which occur abundantly in nature and which have interesting and valuable biological activities. Tarjan and Cheo (Tarjan, A.C., P.C. Cheo [1956] "Nematocidal Value of Some Fatty Acids," Bulletin 332, Contribution 884, Agricultural Experiment Station, University of Rhode Island, Kingston, 41 pp.) report the activity of certain fatty acids against nematodes. In 1977 Sitaramaiah and Singh (Sitaramaiah, K., R.S. Singh [1977] *Indian J. Nematol.* 7:58-65) also examined the response of nematodes to fatty acids. The results of these tests with short chain acids were equivocal, showing nematode-inhibitory action in some instances and stimulatory activity in other instances. Phytotoxicity of these acids was observed at higher concentrations. The short chain fatty acids were also examined by Malik and Jairajpuri (Malik, Z., M.S. Jairajpuri [1977] *Nematol. medit.* 12:73-79), who observed nematode toxicity at high concentrations of the fatty acids.

[0006] Notwithstanding the foregoing (some of the limitations of and problems associated with these approaches are discussed above), there is a need for safe and effective alternatives for controlling nematodes.

[0007] One method for disrupting normal cellular processes is by the use double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). When RNAi corresponding to a sense and antisense sequence of a target mRNA is introduced into a cell, the targeted mRNA is degraded and protein translation of that message is stopped. Although not yet fully understood, the mechanism of this post-transcriptional gene

silencing appears to be at least partially due to the generation of small RNA molecules, about 21 - 25 nucleotides in length, that correspond to the sense and antisense pieces of the RNAi introduced into the cell (Bass, B. L. [2000] "Double-stranded RNA as a template for gene silencing" *Cell* 101:235-238).

[0008] The specificity of this gene silencing mechanism appears to be extremely high, blocking expression only of targeted genes, while leaving other genes unaffected. A recent example of the use of RNAi; to inhibit genetic function in plants used *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* (Chuang, C.-F. and E. M. Meyerowitz [2000] "Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*" *Proc. Natl. Acad. Sci. USA* 97:4985-4990). Chuang *et al.* describe the construction of vectors delivering variable levels of RNAi targeted to each of four genes involved in floral development. Severity of abnormal flower development varied between transgenic lines. For one of the genes, AGAMOUS (AG), a strong correlation existed between declining accumulation of mRNA and increasingly severe phenotypes, suggesting that AG-specific endogenous mRNA is the target of RNAi.

Brief Summary of the Invention

[0009] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences that encode nematode genes, RNAi that selectively targets mRNA transcripts of these essential nematode genes, and methods of using these sequences in nematode control strategies. Such sequences for use according to the subject invention are summarized in Appendix 1. RNAi molecules disclosed herein can be used to inhibit the expression of one or more of these genes in nematodes.

Brief Description of the Drawings

[00010] **Figure 1:** Modular Binary Construct System (MBCS): A series of six, 8-base cutter restriction enzyme sites has been placed between the left and right Ti borders of a previously created kan^R/tet^R binary plasmid.

[00011] **Figure 2:** An exemplary shuttle vector created for cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites.

[00012] **Figure 3:** An exemplary shuttle vector with exemplary inserts.

[00013] **Figure 4:** A suggested RNAi binary vector with exemplary inserts.

[00014] **Figure 5:** Exemplary selectable markers for MBCS.

[00015] **Figure 6:** Exemplary scorable markers for MCBS.

[00016] **Figure 7:** Exemplary RNAi binary vector.

[00017] **Figure 8:** Exemplary RNAi shuttle vector.

Brief Description of the Sequences

[00018] Brief Description of the Sequences can be found in Appendix I.

Detailed Disclosure of the Invention

[00019] The subject invention provides novel methods and compositions for controlling nematodes. More specifically, the subject invention provides polynucleotide sequences and methods of using these sequences in nematode control strategies. A preferred method for controlling nematodes according to the subject invention provides materials and methods for controlling nematodes by using double-stranded interfering RNA (RNAi), or RNA-mediated interference (RNAi). The terms RNAi and RNAi are used interchangeably herein unless otherwise noted.

[00020] In one embodiment of the invention, RNAi molecules are provided which are useful in methods of killing nematodes and/or inhibiting their growth, development, parasitism or reproduction. RNAi molecules of the invention are also useful for the regulation of levels of specific mRNA in nematodes.

[00021] dsRNA (RNAi) typically comprises a polynucleotide sequence identical to a target gene (or fragment thereof) linked directly, or indirectly, to a polynucleotide

sequence complementary to the sequence of the target gene (or fragment thereof). The dsRNA may comprise a polynucleotide linker (stuffer) sequence of sufficient length to allow for the two polynucleotide sequences to fold over and hybridize to each other; however, a linker sequence is not necessary. The linker (stuffer) sequence is designed to separate the antisense and sense strands of RNAi significantly enough to limit the effects of steric hindrances and allow for the formation of dsRNA molecules.

[00022] RNA containing a nucleotide sequence identical to a fragment of the target gene is preferred for inhibition; however, RNA sequences with insertions, deletions, and point mutations relative to the target sequence can also be used for inhibition. Sequence identity may be optimized by sequence comparison and alignment algorithms known in the art (see Gribskov and Devereux, *Sequence Analysis Primer*, Stockton Press, 1991, and references cited therein) and calculating the percent difference between the nucleotide sequences by, for example, the Smith-Waterman algorithm as implemented in the BESTFIT software program using default parameters (e.g., University of Wisconsin Genetic Computing Group). Alternatively, the duplex region of the RNA may be defined functionally as a nucleotide sequence that is capable of hybridizing with a fragment of the target gene transcript.

[00023] As disclosed herein, 100% sequence identity between the RNA and the target gene is not required to practice the present invention. Thus the invention has the advantage of being able to tolerate sequence variations that might be expected due to genetic mutation, strain polymorphism, or evolutionary divergence.

[00024] RNA may be synthesized either *in vivo* or *in vitro*. Endogenous RNA polymerase of the cell may mediate transcription *in vivo*, or cloned RNA polymerase can be used for transcription *in vivo* or *in vitro*. For transcription from a transgene *in vivo* or an expression construct, a regulatory region (e.g., promoter, enhancer, silencer, splice donor and acceptor, polyadenylation) may be used to transcribe the RNA strand (or strands). Inhibition may be targeted by specific transcription in an organ, tissue, or cell type; stimulation of an environmental condition (e.g., infection, stress, temperature, chemical inducers); and/or engineering transcription at a developmental stage or age. The RNA strands may or may not be polyadenylated; the RNA strands may or may not be capable of being translated into a polypeptide by a cell's translational apparatus. RNA

may be chemically or enzymatically synthesized by manual or automated reactions. The RNA may be synthesized by a cellular RNA polymerase or a bacteriophage RNA polymerase (e.g., T3, T7, SP6). The use and production of an expression construct are known in the art (see, for example, WO 97/32016; U.S. Pat. Nos. 5,593,874; 5,698,425; 5,712,135; 5,789,214; and 5,804,693; and the references cited therein). If synthesized chemically or by *in vitro* enzymatic synthesis, the RNA may be purified prior to introduction into the cell. For example, RNA can be purified from a mixture by extraction with a solvent or resin, precipitation, electrophoresis, chromatography, or a combination thereof. Alternatively, the RNA may be used with no or a minimum of purification to avoid losses due to sample processing. The RNA may be dried for storage or dissolved in an aqueous solution. The solution may contain buffers or salts to promote annealing, and/or stabilization of the duplex strands.

[00025] Preferably and most conveniently, RNAi can be targeted to an entire polynucleotide sequence of a gene set forth herein. Preferred RNAi molecules of the instant invention are highly homologous or identical to the polynucleotides summarized in Appendix 1. The homology is preferably greater than 90% and is most preferably greater than 95%.

[00026] Fragments of genes can also be targeted. These fragments are typically in the approximate size range of about 20 nucleotides. Thus, targeted fragments are preferably at least about 15 nucleotides. In certain embodiments, the gene fragment targeted by the RNAi molecule is about 20-25 nucleotides in length. However, other size ranges can also be used. For example, using a *C. elegans* microinjection assay, RNAi "fragments" of about 60 nucleotides with between 95 and 100% identity (to a nematode gene) were determined to cause excellent inhibition.

[00027] Thus, RNAi molecules of the subject invention are not limited to those that are targeted to the full-length polynucleotide or gene. The nematode gene product can be inhibited with a RNAi molecule that is targeted to a portion or fragment of the exemplified polynucleotides; high homology (90-95%) or identity is also preferred, but not necessarily essential, for such applications.

[00028] The polynucleotide sequences identified in Appendix A and shown in the Sequence ID listing are from genes encoding nematode proteins having the functions

shown in Appendix 1. The genes exemplified herein are representative of particular classes of proteins which are preferred targets for disruption according to the subject invention. These classes of proteins include, for example, proteins involved in ribosome assembly; neuro transmitter receptors and ligands; electron transport proteins; metabolic pathway proteins; and protein and polynucleotide production, folding, and processing proteins.

[00029] Genetic regulatory sequences, such as promoters, enhancers, and terminators, can be used in genetic constructs to practice the subject invention. Such constructs themselves can also be used for nematode control. Various constructs can be used to achieve expression in specific plant tissues (by using root specific promoters, for example) and/or to target specific nematode tissues (by using targeting elements or adjacent targeting sequences, for example).

[00030] In a specific embodiment of the subject invention, plant cells, preferably root cells, are genetically modified to produce at least one RNAi that is designed to be taken up by nematodes during feeding to block expression (or the function of) of a target gene. As is known in the art, RNAi can target and reduce (and, in some cases, prevent) the translation of a specific gene product. RNAi can be used to reduce or prevent message translation in any tissue of the nematode because of its ability to cross tissue and cellular boundaries. Thus, RNAi that is contacted with a nematode by soaking, injection, or consumption of a food source will cross tissue and cellular boundaries. RNAi can also be used as an epigenetic factor to prevent the proliferation of subsequent generations of nematodes.

[00031] Nematode polynucleotide sequences disclosed herein demonstrate conserved nucleotide motifs among different nematode genera. Conserved nucleotide motifs strongly suggest that these sequences are associated with viability and/or parasitism and are functionally conserved and expressed in both *Meloidogyne incognita* (root-knot nematode) and *Globodera rostochiensis* and *Globodera pallids* (potato cyst nematodes). The use of these polynucleotides, and RNAi inhibitors thereof, is advantageous because such RNAi can be designed to have broad RNAi specificity and are thus useful for controlling a large number of plant parasitic nematodes *in planta*. Because the genes identified in this disclosure are associated with nematode survival

and/or parasitism, RNAi inhibition of these genes (arising from contacting nematodes with compositions comprising RNAi molecules) prevents and/or reduces parasitic nematode growth, development, and or parasitism.

[00032] Methods of the subject invention include the transformation of plant cells with genes or polynucleotides of the present invention, which can be used to produce nematode inhibitors or RNAi in the plants. In one embodiment, the transformed plant or plant tissue can express RNAi molecules encoded by the gene or polynucleotide sequence introduced into the plant. Other nematode inhibitors contemplated by the invention include antisense molecules specific to the polynucleotide sequences disclosed herein. The transformation of plants with genetic constructs disclosed herein can be accomplished using techniques well known to those skilled in the art and can involve modification of the gene(s) to optimize expression in the plant to be made resistant to nematode infection and infestation. Furthermore, it is known in the art that many tissues of the transgenic plants (such as the roots) can be targeted for transformation.

[00033] RNA-mediated interference (RNAi) of gene expression. Several aspects of root-knot nematode biology make classical genetic studies difficult with this organism. Since root-knot nematodes reproduce by obligatory mitotic parthenogenesis, the opportunity to perform genetic crosses is not available. Microinjection of RNAi can be used to manipulate gene expression in *C. elegans* (Fire, A., S. Xu, M. K. Montgomery, S. A. Kostas, S. E. Driver, and C. C. Mello. [1998] "Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*" *Nature* 391:806- 811). Microinjecting (into adult nematodes) RNAi can turn off specific genes in progeny worms complementary to the coding region of the genes. Moreover, gene inhibition occurs in progeny when RNAi is injected into the body cavity of the adult, indicating the ability of the RNAi to cross cellular boundaries. This RNAi injection method provides a molecular genetic tool that allows for analysis of gene function in root-knot nematodes.

[00034] RNAi can be taken up by *C. elegans* by simply soaking the nematodes in a solution RNAi. This results in targeted inhibition of gene expression in the nematode (Maeda, I., Y. Kohara, M. Yamamoto and A. Sugimoto [1999] "RNAi screening with a non-redundant cDNA set" International Worm Meeting, Madison, WI, abstract 565). Nematodes fed *E. coli* expressing RNAi also demonstrate targeted and

heritable inhibition of gene expression (Sarkissian, M., H. Tabara and C. C. Mello [1999] "A mut-6 screen for RNAi deficient mutants" International Worm Meeting, Madison, WI, abstract 741; Timmons, I. and A. Fire [1998] "Specific interference by ingested dsRNA" *Nature* 395:854; WO 99/32619, hereby incorporated by reference in its entirety).

[00035] Accordingly, one aspect of the instant invention is directed to the control of nematodes comprising contacting nematodes with compositions comprising RNAi molecules specific to the nematode genes disclosed herein. The contacting step may include soaking the nematodes in a solution containing RNAi molecules, feeding nematodes RNAi molecules contained in microbes or plant cells upon which the nematode feeds, or injecting nematodes with RNAi. Nematodes can also be "contacted" and controlled by RNAi expressed in plant tissues that would be consumed, ingested, or frequented by nematodes.

[00036] The RNAi molecules provided to the nematodes may be specific to a single gene. A "cocktail" of RNAi molecules specific to various segments of a single gene can also be used. In addition, a "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) may be applied to the nematodes according to the subject invention.

[00037] In addition to RNAi uptake mediated by transgenic plants, nematodes can be directly transformed with RNAi constructs of cDNAs encoding secretory or other essential proteins to reduce expression of the corresponding gene. The transgenic animals can be assayed for inhibition of gene product using immunoassays or for reduced virulence on a host. Progeny of affected worms can also be assayed by similar methods.

[00038] Procedures that can be used for the preparation and injection of RNAi include those detailed by Fire *et al.*, (1998; <ftp://ciw1.ciwemb.edu>). Root-knot nematodes can be routinely monoxenically cultured on *Arabidopsis thaliana* roots growing on Gamborg's B-5/Gelrite® media. This nematode-host pathosystem is ideally suited for these microinjection experiments since limited root galling results in the parasitic stages (late J2 through adult females) developing outside of the root for easy accessibility for injecting. Another advantage is the parthenogenic reproduction of root-knot nematodes, which makes fertilization by males unnecessary for egg production. The RNAi can be injected into the body cavity of parasitic stages of root-knot nematodes

feeding on *A. thaliana* roots using microinjection. Control nematodes can be injected in parallel with only buffer or an unrelated RNAi. Injected nematodes can be monitored for egg production, and the eggs can be collected for the assays described below. Female root-knot nematodes will typically survive and lay more than 250 eggs following 1 μ l injection of buffer.

[00039] Alternatively, methods are available for microinjecting materials directly into the plant root cells upon which nematodes feed: giant cells or syncytial cells (Böckenhoff, A. and F.M.W. Grundler [1994] "Studies on the nutrient uptake by the beet cyst nematode *Heterodera schachtii* by *in situ* microinjection of fluorescent probes into the feeding structures in *Arabidopsis thaliana*" *Parasitology* 109:249-254). This provides an excellent test system to screen RNAi molecules for efficacy by directly inhibiting growth and development of the nematode feeding upon the microinjected plant cell, or by reducing fecundity and the ability of said nematode to generate pathogenic or viable progeny.

[00040] There are a number of strategies that can be followed to assay for RNAi gene interference. Inhibition of gene expression by RNAi inhibits the accumulation of the corresponding secretory protein in the esophageal gland cells of transgenic J2 hatched from the eggs produced by the injected nematodes. In the first assay, polyclonal antibodies to the target gene product can be used in immunolocalization studies (Hussey, R. S. [1989] "Monoclonal antibodies to secretory granules in esophageal glands of *Meloidogyne* species" *J. Nematol.* 21:392-398; Borgonie, G, E. van Driessche, C. D. Link, D. de Waele, and A. Coomans [1994] "Tissue treatment for whole mount internal lectin staining in the nematodes *Caenorhabditis elegans*, *Panagrolaimus superbus* and *Acrobeloides maximus*" *Histochemistry* 101:379-384) to monitor the synthesis of the target protein in the gland cells of progeny of the injected nematodes, or in any other nematode tissue that fails to express the essential targeted gene. Interference of endogenous gene activity by the RNAi eliminates binding of the antibodies to secretory granules in the glands, or any other target tissue, of the transgenic nematodes, and can be monitored by these *in situ* hybridization experiments. Control nematodes injected only with the injection buffer can be processed similar to the RNAi treated nematodes.

[00041] Another assay is designed to determine the effect of the RNAi on reducing the virulence of J2 progeny of the injected females. Egg masses from injected females can be transferred singly to *A. thaliana* plates to assess the ability of the transgenic J2 to infect roots. The J2 hatching from the eggs transferred to the plates can be monitored; after 25 days the number of galls with egg laying females can be recorded. The *A. thaliana* roots can also be stained with acid fuchsin to enumerate the number of nematodes in the roots. Egg masses from nematodes injected only with the injection buffer can be handled similarly and used as controls. The treatments can be replicated, and the root infection data can be analyzed statistically. These experiments can be used to assess the importance of the target genes in root-knot nematode's virulence or viability. By staining the J2 progeny of the injected females with the antibodies, it can be determined whether RNAi blocks expression of the targeted gene.

[00042] Additional uses of polynucleotides. The polynucleotide sequences exemplified herein can be used in a variety of ways. These polynucleotides can be used in assays for additional polynucleotides and additional homologous genes, and can be used in tracking the quantitative and temporal expression of parasitism genes in nematodes. These polynucleotides can be cloned into microbes for production and isolation of their gene products. Among the many uses of the isolated gene product is the development of additional inhibitors and modifiers. The protein products of the subject polynucleotides can also be used as diagnostic tools. For example, proteins encoded by the parasitism genes, as identified herein, can be used in large scale screenings for additional peptide inhibitors. The use of peptide phage display screening is one method that can be used in this regard. Thus, the subject invention also provides new biotechnological strategies for managing nematodes under sustainable agricultural conditions.

[00043] Antisense technologies can also be used for phytopathogenic nematode control. Antisense technology can be used to interfere with expression of the disclosed endogenous nematode genes. Antisense technology can also be used to alter the components of plants used as targets by the nematodes. For example, the transformation of a plant with the reverse complement of an endogenous gene encoded by a polynucleotide exemplified herein can result in strand co-suppression and gene silencing

or inhibition of a target involved in the nematode infection process. Thus, the subject invention includes transgenic plants (which are preferably made nematode-resistant in this manner, and other organisms including microbes and phages) comprising RNAi or antisense molecules specific to any of the polynucleotides identified herein.

[00044] Polynucleotide probes. DNA possesses a fundamental property called base complementarity. In nature, DNA ordinarily exists in the form of pairs of anti-parallel strands, the bases on each strand projecting from that strand toward the opposite strand. The base adenine (A) on one strand will always be opposed to the base thymine (T) on the other strand, and the base guanine (G) will be opposed to the base cytosine (C). The bases are held in apposition by their ability to hydrogen bond in this specific way. Though each individual bond is relatively weak, the net effect of many adjacent hydrogen bonded bases, together with base stacking effects, is a stable joining of the two complementary strands. These bonds can be broken by treatments such as high pH or high temperature, and these conditions result in the dissociation, or "denaturation," of the two strands. If the DNA is then placed in conditions which make hydrogen bonding of the bases thermodynamically favorable, the DNA strands will anneal, or "hybridize," and reform the original double-stranded DNA. If carried out under appropriate conditions, this hybridization can be highly specific. That is, only strands with a high degree of base complementarity will be able to form stable double-stranded structures. The relationship of the specificity of hybridization to reaction conditions is well known. Thus, hybridization may be used to test whether two pieces of DNA are complementary in their base sequences. It is this hybridization mechanism which facilitates the use of probes of the subject invention to readily detect and characterize DNA sequences of interest.

[00045] The specifically exemplified polynucleotides of the subject invention can themselves be used as probes. Additional polynucleotide sequences can be added to the ends of (or internally in) the exemplified polynucleotide sequences so that polynucleotides that are longer than the exemplified polynucleotides can also be used as probes. Thus, isolated polynucleotides comprising one or more of the exemplified sequences are within the scope of the subject invention. Polynucleotides that have less nucleotides than the exemplified polynucleotides can also be used and are contemplated within the scope of the present invention. For example, for some purposes, it might be

useful to use a conserved sequence from an exemplified polynucleotide wherein the conserved sequence comprises a portion of an exemplified sequence. Thus, polynucleotides of the subject invention can be used to find additional, homologous (wholly or partially) genes.

[00046] Probes of the subject invention may be composed of DNA, RNA, or PNA (peptide nucleic acid). The probe will normally have at least about 10 bases, more usually at least about 17 bases, and may have about 100 bases or more. Longer probes can readily be utilized, and such probes can be, for example, several kilobases in length. The probe sequence is designed to be at least substantially complementary to a portion of a gene encoding a protein of interest. The probe need not have perfect complementarity to the sequence to which it hybridizes. The probes may be labeled utilizing techniques that are well known to those skilled in this art.

[00047] One approach for the use of the subject invention as probes entails first identifying DNA segments that are homologous with the disclosed nucleotide sequences using, for example, Southern blot analysis of a gene bank. Thus, it is possible, without the aid of biological analysis, to know in advance the probable activity of many new polynucleotides, and of the individual gene products expressed by a given polynucleotide. Such an analysis provides a rapid method for identifying commercially valuable compositions.

[00048] One hybridization procedure useful according to the subject invention typically includes the initial steps of isolating the DNA sample of interest and purifying it chemically. Either lysed nematodes or total fractionated nucleic acid isolated from nematodes can be used. Cells can be treated using known techniques to liberate their DNA (and/or RNA). The DNA sample can be cut into pieces with an appropriate restriction enzyme. The pieces can be separated by size through electrophoresis in a gel, usually agarose or acrylamide. The pieces of interest can be transferred to an immobilizing membrane.

[00049] The particular hybridization technique is not essential to the subject invention. As improvements are made in hybridization techniques, they can be readily applied.

[00050] The probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs. Thereafter, the membrane is washed free of extraneous materials, leaving the sample and bound probe molecules typically detected and quantified by autoradiography and/or liquid scintillation counting. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong non-covalent bond between the two molecules, it can be reasonably assumed that the probe and sample are essentially identical or very similar. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred.

[00051] In the use of the nucleotide segments as probes, the particular probe is labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include ^{32}P , ^{35}S , or the like. Non-radioactive labels include, for example, ligands such as biotin or thyroxine, as well as enzymes such as hydrolases or peroxidases, or the various chemiluminescers such as luciferin, or fluorescent compounds like fluorescein and its derivatives. In addition, the probes can be made inherently fluorescent as described in International Application No. WO 93/16094.

[00052] Various degrees of stringency of hybridization can be employed. The more stringent the conditions, the greater the complementarity that is required for duplex formation. Stringency can be controlled by temperature, probe concentration, probe length, ionic strength, time, and the like. Preferably, hybridization is conducted under moderate to high stringency conditions by techniques well known in the art, as described, for example, in Keller, G.H., M.M. Manak (1987) *DNA Probes*, Stockton Press, New York, NY., pp. 169-170.

[00053] As used herein "moderate to high stringency" conditions for hybridization refers to conditions that achieve the same, or about the same, degree of specificity of hybridization as the conditions "as described herein." Examples of moderate to high stringency conditions are provided herein. Specifically, hybridization of immobilized DNA on Southern blots with ^{32}P -labeled gene-specific probes was performed using standard methods (Maniatis *et al.*). In general, hybridization and subsequent washes were carried out under moderate to high stringency conditions that

allowed for detection of target sequences with homology to sequences exemplified herein. For double-stranded DNA gene probes, hybridization was carried out overnight at 20-25° C below the melting temperature (T_m) of the DNA hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature is described by the following formula from Beltz *et al.* (1983):

[00054] $T_m = 81.5^\circ\text{C} + 16.6 \log[\text{Na}^+] + 0.41(\%G+C) - 0.61(\%\text{formamide}) - 600/\text{length of duplex in base pairs.}$

Washes are typically carried out as follows:

- (1) Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).
- (2) Once at $T_m - 20^\circ\text{C}$ for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

[00055] For oligonucleotide probes, hybridization was carried out overnight at 10-20° C below the melting temperature (T_m) of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/ml denatured DNA. T_m for oligonucleotide probes was determined by the following formula from Suggs *et al.* (1981):

[00056] $T_m (^\circ\text{C}) = 2(\text{number T/A base pairs}) + 4(\text{number G/C base pairs})$

[00057] Washes were typically carried out as follows:

- [00058] (1) Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).
- [00059] (2) Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

[00060] In general, salt and/or temperature can be altered to change stringency. With a labeled DNA fragment of greater than about 70 or so bases in length, the following conditions can be used:

Low:	1 or 2X SSPE, room temperature
Low:	1 or 2X SSPE, 42° C
Moderate:	0.2X or 1X SSPE, 65° C
High:	0.1X SSPE, 65° C.

[00061] Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid, and, as noted above, a certain degree of mismatch

can be tolerated. Therefore, polynucleotide sequences of the subject invention include mutations (both single and multiple), deletions, and insertions in the described sequences, and combinations thereof, wherein said mutations, insertions, and deletions permit formation of stable hybrids with a target polynucleotide of interest. Mutations, insertions, and deletions can be produced in a given polynucleotide sequence using standard methods known in the art. Other methods may become known in the future.

[00062] The mutational, insertional, and deletional variants of the polynucleotide sequences of the invention can be used in the same manner as the exemplified polynucleotide sequences so long as the variants have substantial sequence similarity with the original sequence. As used herein, substantial sequence similarity refers to the extent of nucleotide similarity that is sufficient to enable the variant polynucleotide to function in the same capacity as the original sequence. Preferably, this similarity is greater than 50%; more preferably, this similarity is greater than 75%; and most preferably, this similarity is greater than 90%. The degree of similarity needed for the variant to function in its intended capacity will depend upon the intended use of the sequence. It is well within the skill of a person trained in this art to make mutational, insertional, and deletional mutations that are designed to improve the function of the sequence or otherwise provide a methodological advantage.

[00063] PCR technology. Polymerase Chain Reaction (PCR) is a repetitive, enzymatic, primed synthesis of a nucleic acid sequence. This procedure is well known and commonly used by those skilled in this art (see U.S. Patent Nos. 4,683,195; 4,683,202; and 4,800,159; Saiki *et al.*, 1985). PCR is based on the enzymatic amplification of a DNA fragment of interest that is flanked by two oligonucleotide primers that hybridize to opposite strands of the target sequence. The primers are oriented with the 3' ends pointing towards each other. Repeated cycles of heat denaturation of the template, annealing of the primers to their complementary sequences, and extension of the annealed primers with a DNA polymerase result in the amplification of the segment defined by the 5' ends of the PCR primers. Since the extension product of each primer can serve as a template for the other primer, each cycle essentially doubles the amount of DNA fragment produced in the previous cycle. This results in the exponential accumulation of the specific target fragment, up to several million-fold in a

few hours. By using a thermostable DNA polymerase such as *Taq* polymerase, which is isolated from the thermophilic bacterium *Thermus aquaticus*, the amplification process can be completely automated. Other enzymes that can be used are known to those skilled in the art.

[00064] The polynucleotide sequences of the subject invention (and portions thereof such as conserved regions and portions that serve to distinguish these sequences from previously-known sequences) can be used as, and/or used in the design of, primers for PCR amplification. In performing PCR amplification, a certain degree of mismatch can be tolerated between primer and template. Therefore, mutations, deletions, and insertions (especially additions of nucleotides to the 5' end) of the exemplified polynucleotides can be used in this manner. Mutations, insertions and deletions can be produced in a given primer by methods known to an ordinarily skilled artisan.

[00065] The polynucleotide sequences of the instant invention may be "operably linked" to regulatory sequences such as promoters and enhancers. Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is "operably linked" to DNA encoding a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is "operably linked" to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is "operably linked" to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[00066] Polynucleotides and proteins. Polynucleotides of the subject invention can be defined according to several parameters. One characteristic is the biological activity of the protein products as identified herein. The proteins and genes of the subject invention can be further defined by their amino acid and nucleotide sequences. The sequences of the molecules can be defined in terms of homology to certain exemplified sequences as well as in terms of the ability to hybridize with, or be amplified by, certain

exemplified probes and primers. Additional primers and probes can readily be constructed by those skilled in the art such that alternate polynucleotide sequences encoding the same amino acid sequences can be used to identify and/or characterize additional genes. The proteins of the subject invention can also be identified based on their immunoreactivity with certain antibodies.

[00067] The polynucleotides and proteins of the subject invention include portions, fragments, variants, and mutants of the full-length sequences as well as fusions and chimerics, so long as the encoded protein retains the characteristic biological activity of the proteins identified herein. As used herein, the terms "variants" or "variations" of genes refer to nucleotide sequences that encode the same proteins or which encode equivalent proteins having equivalent biological activity. As used herein, the term "equivalent proteins" refers to proteins having the same or essentially the same biological activity as the exemplified proteins.

[00068] It will be apparent to a person skilled in this art that genes within the scope of the subject invention can be identified and obtained through several means. The specific genes exemplified herein may be obtained from root-knot nematodes. Genes, or portions or variants thereof, may also be artificially synthesized by, for example, a gene synthesizer.

[00069] Variations of genes may be readily constructed using standard techniques such as site-directed mutagenesis and other methods of making point mutations and by DNA shuffling, for example. In addition, gene and protein fragments can be made using commercially available exonucleases, endonucleases, and proteases according to standard procedures. For example, enzymes such as *Bal31* can be used to systematically cut off nucleotides from the ends of genes. In addition, genes that encode fragments may be obtained using a variety of restriction enzymes. Proteases may be used to directly obtain active fragments of these proteins. Of course, molecular techniques for cloning polynucleotides and producing gene constructs of interest are also well known in the art. *In vitro* evaluation techniques, such as MAXYGEN's "Molecular Breeding" can also be applied to practice the subject invention.

[00070] Other molecular techniques can also be applied using the teachings provided herein. For example, antibodies raised against proteins encoded by

polynucleotides disclosed herein can be used to identify and isolate proteins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the proteins that are conserved and most distinct from other proteins. These antibodies can then be used to specifically identify equivalent proteins by immunoprecipitation, enzyme linked immunosorbent assay (ELISA), or Western blotting. Antibodies to proteins encoded by polynucleotides disclosed herein, or to equivalent proteins, can readily be prepared using standard procedures known in the art. The genes that encode these proteins can be obtained from various organisms.

[00071] Because of the redundancy of the genetic code, a variety of different DNA sequences can encode the amino acid sequences encoded by the polynucleotide sequences disclosed herein. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding proteins having the same, or essentially the same, amino acid sequence. These variant DNA sequences are within the scope of the subject invention. As used herein, reference to "essentially the same" sequence refers to sequences that have amino acid substitutions, deletions, additions, or insertions that do not materially affect biological activity. Fragments retaining the characteristic biological activity are also included in this definition.

[00072] A further method for identifying genes and polynucleotides (and the proteins encoded thereby) of the subject invention is through the use of oligonucleotide probes. Probes provide a rapid method for identifying genes of the subject invention. The nucleotide segments that are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures.

[00073] The subject invention comprises variant or equivalent proteins (and nucleotide sequences coding for equivalent proteins or for inhibitors of the genes encoding such proteins) having the same or similar biological activity of inhibitors or proteins encoded by the exemplified polynucleotides. Equivalent proteins will have amino acid similarity with an exemplified protein (or peptide). The amino acid and/or nucleotide identity will typically be greater than 60%. Preferably, the identity will be greater than 75%. More preferably, the identity will be greater than 80%, and even more preferably greater than 90%. Most preferably, the identity will be greater than 95%. RNAi molecules will also have corresponding identities in these preferred ranges. These

identities are as determined using standard alignment techniques for determining amino acid and/or nucleotide identity. The identity/similarity will be highest in critical regions of the protein or gene including those regions that account for biological activity or that are involved in the determination of three-dimensional configuration that is ultimately responsible for the biological activity. In this regard, certain amino acid substitutions are acceptable and can be expected if these substitutions are in regions which are not critical to activity or are conservative amino acid substitutions which do not affect the three-dimensional configuration of the molecule. For example, amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type fall within the scope of the subject invention so long as the substitution does not materially alter the biological activity of the compound. Below is a list of examples of amino acids belonging to various classes

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

[00074] In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not detract from the ability to manage nematode-caused diseases.

[00075] An "isolated" or "substantially pure" nucleic acid molecule or polynucleotide is a polynucleotide that is substantially separated from other polynucleotide sequences which naturally accompany a nucleic acid molecule. The term embraces a polynucleotide sequence which was removed from its naturally occurring environment by the hand of man. This includes recombinant or cloned DNA isolates,

chemically synthesized analogues and analogues biologically synthesized by heterologous systems. An "isolated" or "purified" protein, likewise, is a protein removed from its naturally occurring environment.

[00076] Recombinant hosts. The genes, antisense, and RNAi polynucleotides within the scope of the present invention can be introduced into a wide variety of microbial or plant hosts. Plant cells can be transformed (made recombinant) in this manner. Microbes, for example, can also be used in the application of RNAi molecules of the subject invention in view of the fact that microbes are a food source for nematodes

[00077] There are many methods for introducing a heterologous gene or polynucleotide into a host cell or cells under conditions that allow for stable maintenance and expression of the gene or polynucleotide. These methods are well known to those skilled in the art. Synthetic genes, such as, for example, those genes modified to enhance expression in a heterologous host (such as by preferred codon usage or by the use of adjoining, downstream, or upstream enhancers) that are functionally equivalent to the genes (and which encode equivalent proteins) can also be used to transform hosts. Methods for the production of synthetic genes are known in the art.

[00078] Where the gene or polynucleotide of interest is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, certain host microbes are preferred. Certain microorganism hosts are known to occupy the phytosphere, phylloplane, phyllosphere, rhizosphere, and/or rhizoplane of one or more crops of interest. These microorganisms can be selected so as to be capable of successfully competing in the particular environment (crop and other habitats) with the wild-type microorganisms, provide for stable maintenance and expression of the gene expressing a polypeptide of interest, and, desirably, provide for improved protection of the protein/peptide from environmental degradation and inactivation.

[00079] A large number of microorganisms is known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, *e.g.*, genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*,

Arthrobacter, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, *e.g.*, genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are the pigmented microorganisms.

[00080] Methods of the subject invention also include the transformation of plants or plant tissue with genes which encode the RNAi molecules of the present invention. In one embodiment, the transformed plant or plant tissue expresses antisense RNA and/or RNAi. Transformation of cells can be made by those skilled in the art using standard techniques. Materials necessary for these transformations are disclosed herein or are otherwise readily available to the skilled artisan.

[00081] Additional methods and formulations for control of pests. Control of nematode pests using the RNAi molecules of the instant invention can be accomplished by a variety of additional methods that would be apparent to those skilled in the art having the benefit of the subject disclosure. A "cocktail" of two or more RNAi molecules can be used to disrupt one or more of the genes identified herein. The "cocktail" of RNAi molecules may be specific to segments of a single gene or the entire gene. A "multigene cocktail" of RNAi molecules specific to two or more genes (or segments thereof) is also encompassed by the instant invention. In another embodiment of the instant invention, the disclosed RNAi molecules, cocktails, and/or multigene cocktails thereof, may be used in conjunction with other known nematode control agents and methodologies. Such cocktails can be used to combat the development of resistance by nematodes to a certain inhibitor or inhibitors.

[00082] Compositions of the subject invention which comprise RNAi molecules and carriers can be applied, themselves, directly or indirectly, to locations frequented by, or expected to be frequented by, nematodes. Microbial hosts which were transformed with polynucleotides that encode RNAi molecules, express said RNAi molecules, and which colonize roots (*e.g.*, *Pseudomonas*, *Bacillus*, and other genera) can be applied to the sites of the pest, where they will proliferate and be ingested. The result is control of the pest. Thus, methods of the subject invention include, for example, the application of recombinant microbes to the pests (or their locations). The recombinant microbes may also be transformed with more than one RNAi molecule thereby delivering a "cocktail" of RNAi molecules to the nematode pests. A carrier may be any substance suitable for

delivering the RNAi molecules to the nematode. Acceptable carriers are well known in the art and also are commercially available. For example, such acceptable carriers are described in E.W. Martin's *Remington's Pharmaceutical Science*, Mack Publishing Company, Easton, PA.

[00083] All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety to the extent they are not inconsistent with the explicit teachings of this specification.

[00084] Following are examples that illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

Example 1— Production of Hairy Roots for RNAi Testing

[00085] A hairy root assay system was developed for testing the anti-nematode activity of RNAi molecules.

[00086] *Agrobacterium rhizogenes*: Several *Agrobacterium rhizogenes* strains produce hairy roots on a variety of plant species. *A. rhizogenes* strains, A4, 15834, 8196 and LBA4404 demonstrate hairy root development on tomato and sugar beet, with A4 being the most efficient. The *A. rhizogenes* strain K599 demonstrated very efficient formation on transgenic soybean hairy roots and was also effective on sugar beet and *Arabidopsis*. However, stain K599 failed to produce hairy roots on tomato tissues possibly due to hyper-virulence.

[00087] Hairy root production: Transgenic hairy roots were identified by stable GUS expression in tomato, sugar beet, soybean and *Arabidopsis*. The construct pAKK1401 (pNOS / NPT-II / tNOS // pSU / GUS / tNOS) was used to produce hairy roots when transformed into *A. rhizogenes* strains A4 or K599. Transgenic roots were identified by GUS expression.

Example 2 – Protocol for Electro-competent *Agrobacterium* and Electroporation

[00088] Electro-competent *Agrobacterium* Protocol:

- [00089] 1. Grow *Agrobacterium* overnight in 5 mls LB + antibiotics at 30°C on shaker (for *Agrobacterium rhizogenes* strain K599 no antibiotics are needed).
- [00090] 2. Use the 5 mls of overnight culture to inoculate 500 mls LB + antibiotics at 30°C on shaker. Grow overnight.
- [00091] 3. Add liquid culture in eight 50 ml polypropylene orange cap tubes.
- [00092] 4. Centrifuge 10 min., 4000 rpm, 4°C.
- [00093] 5. Resuspend cells in each tube with 20 mls 10% glycerol (on ice)
- [00094] 6. Centrifuge 10 min., 4000 rpm, 4°C.
- [00095] 7. Resuspend cells in each tube with 10 mls 10% glycerol (on ice).
- [00096] 8. Centrifuge 10 min., 4000 rpm, 4°C.
- [00097] 9. Resuspend cells in each tube with 2 mls 10% glycerol (on ice).
- [00098] 10. Aliquot 50 µl into cold Eppendorf tube and place onto dry ice.
- [00099] 11. Store electro-competent cells at -80°C. These cells can be used for up to two years.

[000100] Electroporations:

- [000101] 1. Add 1 µl to 5 µl of DNA (resuspended in H₂O and not TE or other buffer) to 50 µl of *Agrobacterium* electrocompetent cells and mix.
- [000102] 2. Transfer 20 µl of DNA/*Agrobacterium* mix to cuvette.
- [000103] 3. Electroporate:
25µF, 400 Ω resistance, 2.5 volts (0.2cm cuvette) or 1.8 volts (0.1cm cuvette for BioRad electroporator. 330 µF, 4000 kΩ, low w, fast charge rate for BRL Electroporator.
- [000104] 4. Add 1ml of LB and transfer to Eppendorf tube.
- [000105] 5. Shake at 30°C for 2 hours.
- [000106] 6. Centrifuge down cells (2 min. 14 krpm).
- [000107] 7. Plate all onto LB + antibiotics (most *Agrobacterium* strains are naturally streptomycin resistant).

Example 3 – Protocol for Production of Transgenic Hairy Roots on Soybean

[000108] Seed Sterilization. Rinse the soybean seed with 70% ETOH for 2-5 min. Remove and add 20% Clorox and shake for 20-25 min. Rinse 3X with sterile water. Plate the seed, 5 seed per plate, onto ½ MSB5 + 2% sucrose + 0.2% gel (referred to as ½ MSB5). Place seed into chamber at 25°C, 16/8 photoperiod for 5-7 day (depending on genotype) germination period. After 1 week seedlings can be placed into cold room for longer storage if necessary (not to exceed 2 weeks).

[000109] Agrobacterium Preparation. For *Agrobacterium rhizogenes* strain K599, take a small sample from frozen glycerol into 25-50 ml of NZYM media with 50 mg/L kanamycin in a 125-250 ml Erlenmeyer flask. Place onto shaker at 28-30 °C for 16 - 20 hours. Pour sample into centrifuge tube and centrifuge the bacterium at 4000 rpm for 10 min. Pour off supernatant and re-suspend the pellet with an equal volume of liquid ½ MSB5 + 200 µM acetosyringone. Use pipette to re-suspend the pellet and homogenize the sample (remove all clumps). To determine O.D., prepare a 1:10 dilution by putting 900 µl ½ MSB5 into cuvette and add 100 µl of bacterial sample. Determine the O.D.₆₆₀ and calculate the volume needed to adjust (dilute) OD to approximately 0.2 for inoculation. Check final O.D.

[000110] Explant Preparation and inoculation. Place a sterile filter paper onto plates of 1/2 MSB5. Cut soybean cotyledons just above the shoot apex and place onto plate. Lightly scar the cotyledon's abaxial surface (flat side, upper surface that reaches toward sun) with a scalpel blade. Cut each cotyledon transversely into 2-3 pieces (no smaller than 1 cm). Add approximately 10 ml of prepared bacterial solution to each plate and allow cotyledons to incubate for 1 hr. Remove the bacteria using a vacuum aspirator fitted with sterile pipette tip, ensure that there is no standing liquid. Orient all explants with abaxial surface up and wrap plates for a 3 day co-culture, 25°C in light (16/8 photoperiod).

[000111] Hairy root selection and maintenance. After 3 day co-culture, wash explants with liquid ½ MSB5 + 500 mg/L carbenicillin. Transfer the explants abaxial side up to selection media, ½ MSB5 supplemented with 500 mg/L carbenicillin and 200 mg/L kanamycin. Roots should develop in approximately 2-3 weeks. The roots will form primarily from the cut vascular bundles with other roots developing from the small cuts on cotyledon surface. Remove roots (>1cm in length) and place onto replica media with

transfers to fresh media every 2 weeks to prevent *Agrobacterium* overgrowth. After 6-8 weeks on selection the roots can be moved to media without kanamycin, however carbenicillin must remain in media for several months for continued suppression of *Agrobacterium*. At this stage roots can be used for testing RNAi for nematode control. Sterilized nematodes can be added and observed for RNAi affects.

Example 4 – Testing of RNAi for Plant Parasitic Nematode Control.

[000112] Various types of nematodes can be used in appropriate bioassays. For example, *Caenorhabditis elegans*, a bacterial feeding nematode, and plant parasitic nematodes can be used for bioassay purposes. Examples of plant parasitic nematodes include a migratory endo-parasite, *Pratylenchus scribneri* (lesion), and two sedentary endo-parasites, *Meloidogyne javanica* (root-knot) and *Heterodera schachtii* (cyst).

[000113] *C. elegans*: RNAi vectors can be tested through expression of the RNAi in *E. coli*. *C. elegans* are fed *E. coli* and assayed for their growth by measuring growth of nematodes, production of eggs and viability of offspring. Another approach is to inject dsRNA directly into living nematodes. Finally, soaking nematodes in a solution of *in vitro*-prepared RNAi can quickly establish efficacy of treatment.

[000114] *P. scribneri*: The *P. scribneri in vitro* feeding assay uses a corn root exudate (CRE) as a feeding stimulus and both the red dye Amaranth or potassium, arsenate as feeding indicators. Feeding is confirmed after seven days by the presence of red stained intestinal cells in live worms exposed to the Amaranth or death of worms exposed to arsenate. This bioassay is used to test soluble toxins or RNAi. *P. scribneri* has also been cultured on wild type roots of corn, rice and *Arabidopsis*, and on A. rhizogenes-induced hairy roots of sugar beet and tomato. *P. scribneri* is very valuable in evaluating transgenic hairy roots because of the non-specific feeding of these worms.

[000115] *M. javanica*: Nematode eggs are sterilized using bleach and are used to inoculate hairy roots expressing RNAi. Nematodes are assessed for their growth by measuring knots, egg masses or production of viable eggs. An alternative approach is to microinject dsRNA directly into root feeding sites or into living female nematodes.

[000116] *H. schachtii*: Cultures of this nematode were maintained on sugar beets. Nematodes eggs are sterilized using bleach and used to inoculate hairy roots

expressing RNAi. Nematodes can be assessed for their growth by measuring knots, egg masses or production of viable eggs.

Example 5 – Plant Expression Vectors for RNAi

[000117] Modular Binary Construct System (MBCS): An important aspect of the subject disclosure is the Modular Binary Construct System. The MBCS eases the burden of construct development by creating modular pieces of DNA that can be easily added, removed, or replaced with the use of low frequency cutting restriction enzymes (8-base cutters). These constructs are useful for delivery of a variety of genes to plant cells and is not limited to the delivery of RNAi genes. To develop this system, a series of six, 8-base cutter restriction enzyme sites was placed between the left and right Ti borders of a previously created kan^R/tet^R binary plasmid (Figure 1). The production of both kan^R and tet^R MCBS aids the testing of constructs using different strains of *Agrobacterium rhizogenes* in different plant species. In addition to the MBCS, a series of shuttle vectors were created that aid in the cloning of useful DNA fragments by containing the multi-cloning site (MCS) of a modified Bluescript plasmid flanked by 8-base restriction sites (Figure 2). With six 8-base cutter sites, each site is, preferably, reserved for a particular function (Figures 3 and 4). Because of the close proximity of the *Pme* I and *Sgf* I sites to the left and right border of the binary vector, these sites are, preferably, reserved for gene tagging and enhancer trap experiments. The *Not* I site is, preferably, reserved for plant selectable markers (Figure 5). The *Pac* I site is reserved, preferably, for Plant Scorable Markers (Figure 6). The *Asc* I site is, preferably, reserved for RNAi experiments (Figures 7 and 8), while the *Sbf* I site is, preferably, reserved for anti-nematode proteins. The restriction sites that are denoted in the Figures are, preferably, reserved for the denoted insertions; however, the MCBS binary and shuttle vectors do not require the restriction sites to contain these suggested inserts.

[000118] Plant Selectable Markers for MBCS: To further develop the MBCS, a series of plant selectable markers were added to the MBCS (Figure 5). Plant selectable markers that were added to the MBCS include: pNOS/NPT-II/tNOS (kan^R), pNOS/Bar/tNOS (basta^R for dicots), pUBI/Intron-Bar/tNOS (basta^R for monocots), and pUBI/Intron-PMI/tNOS (mannitol isomerase^R).

[000119] Reporter Genes for MBCS: Four exemplary reporter genes are used in the MBCS are provided in Figure 6 and Appendix 2. GUS, a nuclear localized GUS, GEP, and the anthocyanin transcriptional activator *papIC* genes into the MBCS.

[000120] Promoters for MBCS: We cloned several useful constitutive and nematode-inducible promoters (Figures 6, 7 and Appendix 2). Constitutive promoters include the SuperUbiquitin promoter from pine (pSU) and two promoter regions from the Strawberry Banding Vein virus (pSBV₁ and pSBV₂). Seven nematode-inducible promoters from *Arabidopsis* were also been cloned.

[000121] The following Scorable marker clones have been constructed and placed in the MBCS, NPT-II binary vector (pNOS/NPT-II/tNOS):

Intron/GUS/tNos	Intron/NLS-GUS/tNOS	Intron/GFP/tNOS
pSU/Intron/GUS/tNOS	pSU/Intron/NLS-GUS/tNOS	pSU/Intron/GFP/tNOS
pSBV ₁ /Intron/GUS/tNOS	pSBV ₁ /Intron/NLS-GUS/tNOS	pSBV ₁ /Intron/GFP/tNOS
pSBV ₂ /Intron/GUS/tNOS	pSBV ₂ /Intron/NLS-GUS/tNOS	pSBV ₂ /Intron/GFP/tNOS
pKT/Intron/GFP/tNOS		
pKA/Intron/GFP/tNOS		

Example 6 — Control of Plant parasitic nematodes using RNAi *in planta*

[000122] Production of RNAi Vector. The RNAi shuttle vector to be used is adapted from the Modular Binary Construct System (MBCS - See Example 5). RNAi shuttle vectors preferably comprise a promoter, intron, antisense RNAi, stuffer fragment, sense RNAi, and terminator (See Figures 7 and 8 and Appendix 2 for more details). The plant promoter can be constitutive, tissue-specific or nematode-inducible. The intron is necessary to eliminate expression in *Agrobacterium*.

[000123] The anti-sense and sense RNAi molecules comprise nematode-specific sequences and are disclosed herein. These genes are associated with pathogenesis, growth, or other cellular function in nematodes. An exemplary group of RNAi sequences for use in plant/nematode control may be based upon:

[000124] 1. Genes specific for nematode esophageal gland cells.

[000125] 2. Genes specific for plant parasitic nematodes but not other free living nematodes.

- [000126] 3. Genes common to all plant parasitic nematodes.
- [000127] 4. Genes common to all nematodes (nematode-specific).
- [000128] 5. Genes specific for important tissues or cell types.
- [000129] 6. Genes from large gene families.
- [000130] 7. Genes involved in nematode signal transduction or other cellular pathways.

[000131] Appropriate RNAi constructs allow for the formation of dsRNA molecules (the sense and antisense strands join to form the dsRNA). The terminator sequence adds a poly-A tail for transcriptional termination. The RNAi shuttle vector can then be subcloned into the MBCS and transformed into *Agrobacterium rhizogenes*.

[000132] Plant Transformation with RNAi Vectors. An exemplary transformation system for generating hairy roots using *Agrobacterium rhizogenes* is provided below. The RNAi vector once introduced into the MBCS can subsequently (as a binary vector) be transformed in *A. rhizogenes* using, for example, the electroporation protocol of Example 2. Once the *A. rhizogenes* is confirmed to contain the plasmid, it is then used in generating hairy roots (See Example 3). Using this protocol transgenic hairy roots expressing RNAi are isolated, cultured and tested.

[000133] Testing of RNAi Vector for Nematode or Plant Pathogen Resistance. RNAi expressing hairy roots can be inoculated with sterilized nematodes. Infested hairy roots can be observed and the effect on nematodes determined. An alternative approach involves the microinjection of RNAi directly into root feeding sites (giant-cells for root-knot nematode, and syncytia for cyst nematodes) or into living female nematodes.

Example 7 – Insertion of Genes Into Plants

[000134] One aspect of the subject invention is the transformation of plants with genes encoding proteins of the present invention. Transformation of plants as described herein can be used to improve the resistance of these plants to attack by the target pest.

[000135] Genes, polynucleotides, and/or RNAi molecules as disclosed or suggested herein can be inserted into plant cells using a variety of techniques which are

well known in the art. For example, a large number of cloning vectors, for example, pBR322, pUC series, M13mp series, pACYC184, pMON, *etc.*, are available for preparation for the insertion of foreign genes into higher plants via injection, biolistics (microparticle bombardment), *Agrobacterium tumefaciens*, or *Agrobacterium rhizogenes*-mediated transformation, or electroporation as well as other possible methods. Once the inserted DNA has been integrated into the genome, the genetically modified-cell(s) can be screened via a vector carried-selectable marker that confers on the transformed plant cells resistance to a biocide or an antibiotic, such as kanamycin, G418, bleomycin, hygromycin, chloramphenicol, or bialophos, *inter alia*. The transformed cell will be regenerated into a morphologically normal plant. The transgene(s) in the transgenic plant is relatively stable and can be inherited by progeny plants.

[000136] If a transformation event involves a germ line cell, then the inserted DNA an corresponding phenotypic trait(s) will be transmitted to progeny plants. Such plants can be grown in the normal manner and crossed with plants that have the same transformed hereditary factors or other hereditary factors. The resulting hybrid individuals have the corresponding phenotypic properties.

[000137] It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

We claim:

1. An RNAi molecule, optionally comprising a linker, wherein at least one strand of said RNAi is encoded by a DNA sequence selected from the group consisting of SEQ ID NO: 1 through SEQ ID NO: 139.

2. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 1.

3. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 2.

4. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 3.

5. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 4.

6. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 5.

7. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 6.

8. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 7.

9. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 8.

10. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 9.

10. 11. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
10.
11. 12. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
11.
12. 13. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
12.
13. 14. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
13.
14. 15. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
14.
15. 16. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
15.
16. 17. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
16.
17. 18. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
17.
18. 19. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
18.
19. 20. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
19.
20. 21. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
20.

21. 22. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 21.
22. 23. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 22.
23. 24. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 23.
24. 25. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 24.
25. 26. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 25.
26. 27. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 26.
27. 28. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 27.
28. 29. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 28.
29. 30. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 29.
30. 31. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 30.
31. 32. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 31.

34

32. 33. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

33. 34. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

34. 35. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

35. 36. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

36. 37. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

37. 38. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

38. 39. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

39. 40. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

40. 41. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

41. 42. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

42. 43. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

43. 44. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
43.
44. 45. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
44.
45. 46. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
45.
46. 47. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
46.
47. 48. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
47.
48. 49. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
48.
49. 50. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
49.
50. 51. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
50.
51. 52. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
51.
52. 53. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
52.
53. 54. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
53.

54. 55. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
54.
55. 56. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
55.
56. 57. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
56.
57. 58. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
57.
58. 59. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
58.
59. 60. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
59.
60. 61. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
60.
61. 62. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
61.
62. 63. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
62.
63. 64. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
63.
64. 65. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
64.

65. 66. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 65.
66. 67. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 66.
67. 68. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 67.
68. 69. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 68.
69. 70. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 69.
70. 71. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 70.
71. 72. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 71.
72. 73. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 72.
73. 74. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 73.
74. 75. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 74.
75. 76. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 75.

76. 77. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
77. 78. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
78. 79. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
79. 80. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
80. 81. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
81. 82. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
82. 83. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
83. 84. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
84. 85. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
85. 86. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
86. 87. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:

88. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
87.
89. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
88.
90. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
89.
91. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
90.
92. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
91.
93. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
92.
94. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
93.
95. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
94.
96. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
95.
97. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
96.
98. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO:
97.

99. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 98.

100. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 99.

101. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 100.

102. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 101.

103. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 102.

104. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 103.

105. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 104.

106. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 105.

107. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 106.

108. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 107.

109. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 108.

110. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 109.

111. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 110.

112. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 111.

113. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 112.

114. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 113.

115. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 114.

116. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 115.

117. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 116.

118. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 117.

119. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 118.

120. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 119.

121. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 120.

122. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 121.

123. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 122.

124. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 123.

125. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 124.

126. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 125.

127. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 126.

128. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 127.

129. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 128.

130. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 129.

131. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 130.

132. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 131.

133. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 132.

134. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 133.

135. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 134.

136. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 135.

137. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 136.

138. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 137.

139. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 138.

140. An RNAi molecule according to claim 1, wherein said DNA sequence is SEQ ID NO: 139.

141. A transgenic plant or transgenic plant tissue comprising an RNAi molecule according to any of the preceding claims.

142. A method of disrupting cellular processes in a nematode comprising the steps of:
- (a) providing a composition comprising a compound according to any of the preceding claims; and
 - (b) contacting a nematode with said composition.

143. An isolated promoter comprising the following nucleotide sequence:

```
aacagcccaagataaacagaaaagtcaaagggtgttcgaaa
gaccacttgtgactaaggatcatttcatccataattatctggtagca
cagactcatgataaetgcgaggaaacacaagttctttacagtcgattc
aaagacactttctcttttacggtttcattgaaggagccgaccagaat
atgtcagagaagcttttctactgtgggttaatttcattaatctatcca
ggtgaaaacctcaaggagatctctcttctcccaaaagacctctacag
ggcaatcaaaaactacagaaccagagtttgtagtgcacagagtagac
caatctacctgagaatcacgagtaccttcctagagtgggaaaatgat
gacatccttattccataaccactggattgaggtaggactatccaatgg
aaaaattccatgggacaagtcataataagaagaccgcaacagtcgagt
atcttccagagataactgcactcagacctaaaaggataaaaagcagta
tataatcagtgtactaagatcttcgcagattcaaagaagaagcttaa
ctatgctgatgacaagataattctaataagcaattattcagaattaa
tcaaggagaaagaattaataactctttcagaatatgaagcccgcttt
acaagtggccagctagctatcactgaaaagacagcaagacaatggtg
tctcgatgcaccagaaccacatctttgcagcagatgtgaagcagcca
gagtgggtccacaagacgcactcagaaaaggcatcttctaccgacaca
gaaaaagacaaccacagctcatcatccaacatgtagactgtcgttat
gcgtcggctgaagataagactgaccccaggccagcactaaagaagaa
ataatgcaagtggctcctagctccacttttagctttaataattatgttt
cattattattctctgcttttgctctctatataaagagcttgatattt
catttgaaggcagaggcgaacacacacacagaacctccctgcttaca
aaccatgtattgtagctaaacctcttaggag.
```

144. An isolated promoter comprising the following nucleotide sequence:

45

```
tggtggggacaatggatccggtctgcgtagcaacaaggctg
aaaaagattaaacagaaacctgtgatcattagcgttggaccaccacc
aaaacctcctgagccaccaaaagcctccagagcctgaaaaaccaaagc
ctccaccagcacctgaaccaccaaaagcatgtatgcaagccaccttac
tgcaacagttgtgatgttgtgtctgttactacctatgaaagtggaag
cggctgcaccattctttgagtcataatcgcgtagcatagccttcat
gttaagtcctgtatttagccaataactaattcatcatgttctcatgct
tttttggtttatttctttttctcaaatatgaatctctgttgtttgtcc
ctccctgtttataatttagtcgcttctttgacacaagaagtctcatg
agttcatgctaaagaaaaataaaagttcaaattaaaacaccaaagtgtt
tgattaatttccataaacctgtgaagcagaaagttagtcattgttgac
ctgaacagagccttaggaagtccttgaaggacatatcttcaagtgtta
ttgggtcgtagcactcttaggccatttaacttcattgagccatttaa
attatgcaaaacaagaaatgagacatatggaaacattagggttctta
caggaaaaaataggaaaaagcagggacaactaaacaaaaattcagaa
acaagaggcaagtggacgaccacggcgtaagatcaacatgtggtgat
gtgcatgagaccaagaccattttttctcgttcttcaacgcacacttg
gtcttttcttatgtttgttgcatttctttatttaggcagaccctctct
cttttttaatataggatagtaaaaaatatatgattttattttgttgaaa
cattttgaggttaaaacctaaacttatagtaagcattttagtagagtga
tttctatatacgacatctatcaacatgaccttaacaaaaaaatatt
gatgaaactactttaagtagtaaaacctaaagcaattaaaatttctt
ttaaattagtagtttgtgtaaattaatgacatgattgcgtcgaaag
aaatcaaaacagttatatcgtgaacttaggagaatgttttatatcgt
gtttcaacacatgattgctagcatatgtgtagggtgcgtagacgtta
cataacaatcatcactcgtaaatatcaaagtgggttctgagagaaac
aaagggttatgattttcccaactgcactagttgtgtattgtttcttt
cacacgtatgcttctgagttctgcccaaagtggaaattaaagcagag
ttgggagagatcataatttatttaggggttcgttatgctcaagtcattga
cgtaaaatgaaaatttgtttttattctttcaccaacacaaagaatag
ctagttatctctttttttatatataacaattcatgaagttgatcagc
tttatacacatcatccaatcgaattgctaattctagagatggaaatat
caggatagagccaataagatatcaaattcaatggaccattttctcc
atgtgctaattcatacaatctgtttttgtctgctttattttgatgatg
atgctgagcgttttttaagtgtgaactaagatctagctaaccacaaaca
aagatggctctcttctgtctttgtcgtataagagcaagagagtggttt
gattcaatttttaaaattctaaataaaaactccaaccgtgaatccagc
catgaaaactcttttttagaaaatccttttttataacaaataattctct
tgcttcttcttcttcttctgtttatttcaccttttttggtttcttttag
ctcagaaaaagcccattctttttttctattcttgtttattttaatca
tactgtgcgtttctacaaagtttgttcctttcttcttcaactctctc
actcacagtccagagatctgtttctttttcttttttttgctttcactc
ttctcttccagt.
```

145. An isolated promoter comprising the following nucleotide sequence:

agcaaagcaagaacaccagagaagaagaaaagcactacaga
gaaaaatgtgagcttaagcgcctctccaacaacacttctctgggagtc
taaaggatgctgcaaaaagccttggtggtgagacttccgcataatttc
caagcatgggtttatTTTTgttagcacacaaactatctgaccctcga
cttggatTTTTcttctgcagtttgtccaactacattgaaacggatatg
caggcaacatgggatcatgaggtggccatctcgtaagattaacaaag
tgaacagggtcactaaggaaaatacagacgggtactggactcgggtccaa
gggtgtagaaggaggactaaagtctgactcagcaactggcgaattcat
tgcagttagaccttttattcaagaaattgatacccaaaagggctctgt
cgtctcttgataatgatgcacatgcaagaagaagt caggaggatatg
cctgacgatacttcattcaagctccaggaagctaaatctgtcgacaa
tgccattaagttagaggaggatataccatgaatcaagcaagaccag
gtaagaacttctctatccataaaccatagatggagcgttagaatct
taatccatTTTTcagtttttgcaggatcattcatggaggttaatgcta
gtggtcagccatgggcttggtggccaaagagtctggcttgaatggc
agtgaaggaataaagagcgtttgcaacttaagctctgtggaaatttc
agatggaatggatccaacaatccgatgcagtggcagtatgttgaaac
ctaaccaatccatgtcatgcagcatatcagattcatcaaattggctca
ggcgcagttctgcgtggaagctcatctacttccatggaagattggaa
ccaaatgagaaccacacacagtaatagcagcgagagtggatcaacaa
cgctgatcgtaaaaggccagttatagagaagacactgtacgtttcaag
ttcgagccatcagttgggtgtcctcagctctacaaagaagttggaaa
acgttttaaaactgcaggacgggtcgtttcagctgaagtacttggatg
atgaagaagaatgggtgatgctgggttacagattctgatctccaagaa
tgtttggagatattacatgggtatgggaaaacactcgggtgaagtttct
cgttcgtgatttgtctgccccctctaggtagttctgggtggcagtaatg
gttatcttggaacaggcttatgacgtcgtaagacatagacacacaca
gttatgtattcccagtgaaagaatgttggtttatttctctagatatta
gtatgcttataaataggcatgaaggagaaagacaattttgggtatagt
ggagttcagcagaaaatgtatatgtttttcgttttatatgaatcag
agaataaaaagttggatgttatatctacgttgctaattgttgtaacctgc
tcacccatctttcatataagaaaagagaaacacttttagttatccctg
tgatgcagaatcgtattctttgttatctctccattcctgtggaaacc
aaciaaagtcactaaatttcgggttaattgggttggtttttaagtcaa
cgaggacttgatttttagttgggcttgggcctataattgtgttcatca
ttgggttttttcccccttatcagtttaacgtccatatccatatcttt
ttcttttttaacggcgaagttcatatccatatcttatgatgtgcct
aaaagagggagaagatgcgaagacagaattttcatatttgaaaggggt
tcgatatcgatatgtgggaaacgaatcaagggtcaaaaaactcagtcata
atagttgaaatttaaaaaattttattaattcaatccgattgggttcgt
tttggttatgggttcgggtctatatcatcaaaccaatcggtttgggtcct
aaagataattataaatattcaccaacaccagtggttaaacacatatca
acaaacctaaagttagataaacaagaga.

146. An isolated promoter comprising the following nucleotide sequence:

```
aat tgg cact ctt ctt ctt gct ggg ttc caa aag aa ac ga at
ca at at gt gca aca aga ag ag ct cc aga ag cag tca tct tct aa at
ct ta at cta aca ac ag ct ca aga a gaa aaaa at tcc at ag ct ag a
ga ac aca a ag tca ca ag ac ga cgt cgt ag ag gc aca a ag tca a ac ct
ga at gg ct ta ag cc ga act ga gt ggt ttt g act ag acc at cat caga
aa ag tct cca ag ac gg tag tcc gat gtt ag at cgt cca ag ta at ttt
tgg ttt tgt tgg tct ca cgt ttt cag ct gcc at ttt ga ttt cag ttt
ggg ctt ttc ctt at ctt aag gcc aat ttt cat ttt ag gtt tag ttt
at ttt gat cat ttt at ctt act aaaa ag gct tcc gct ttt cga ga aat t
agg gtt tct tct gtt cgt cgt cact cag gtt tgt gc ct ca ac ga c
tg ctt cact tct ag ct tga ttt ctt ctt cgt ttt at at gt ata ct g
ta cat tag at ttt ctt gtt tct cga gct tct gct at ag at ttt gat
tct ttt ttt tgt tgt ctt tgt ttt cgt ttt cc ag gat cag at ctt ag ct
aa at tg ag aca ag ct caa aat ga ggt act tga cgc at ctt ctt ac at
cact gtt ta at tag aga aca at ac gt ctt ctt ga at cgt gat tca ga ga
cgt at tgt tct tct gtt cat at gca at aag ttt aat tag aga aca at a
cgt ctt ctt ga at cgt gat tgt ttt ttt gg at gtt gc gtt at tga tag ct t
tat gat gtt aat ag tct ag gatt ga cac ga ag ttt gtt ctt gc ag ttt t
gc at aa at gct ctt tta cta ag gct cta aat ttt gg at ga caa at cta
aat ctt gc ct ca taaa aat ttt ag gtt gta tta ag ata ag at ttt ttt g
tat gg tag tgt cta at at gtt ggt tgt tcat gtt ga ggt tgt ca at g
ttgt gta ttt ttt gtt ttt ag tta at ttt gct ta act ctt gtt ctt ttt g
tgg gtt aat ac ag ta ag ct tca g ag tga ggc cgt tct gta ag cc at c
act act at ca cag gga aat cc gagg caa aag aa ac gta act ttt gtt cga
gact at tga gct cc ag at cgt ctt ga aga act at ga cc ct caa aag g
aca ag cgt ttt cag tgg at ctt gtt ca ag tt acc ac at at cccc cgt cct
aaa at ga ag at ctt gcat gct cgg ag at gcc ag cat gtt ga ag ag gt
gat at at ctt ttt cat gg aa at tga tcat ttt ttt gtt ctt gtt ttt ttt gtt
ata at ggt ttt ttt gtt ctt ctt ttt ggt ggt ctt at tga ttt ctt
tg at gtt gta at at gtt ctt ctt ga at gta g at gcat gat gtt ttt cgg aa
ttt ggt ctt ttt gtt ttt at ttt ag gct tcat ttt ctt gc at aat ta aat at t
tg ctt at ttt cat ctt gta tct ttt cgt ag gct gaga ag at ggg gtt g
gaaa ac at gg at gtt g ag tct cta aaaa ag ct ta aca aga aca ag aa
act cgt ca aga ag ct tgc aa a gaa at acc at gct ttt ctt tgg cct ctt g
agt ctt gtt catta ag cag at tct cgt ctt ctt tgg tct ggt ctt aac
aagg cagg ca ag ttt ctt gg ct ac ag cta at at tcc at tgt tct ttt ttt
ac at cc gtt ttt gat ttt tgg at ag gtt ttt ag tag tct at ttt ctt ttt gtt
ca at gtt ctt ttt gata ca at gcc aat ctt ttt at ctt gtt ga gatt at g
ctt ctt ttt gat gat tct ta ag ta ac at tct ttt gct ttt act ttt ac aca
gg aaa at tcc ca act ctt gtt ga gcc acc ag ga at cct tgg ag tca aa
ggt ga at gaa aca aag gca ac ag tga ag ttt cc ag ct ga aga ag gtt c
tgt gcat ggg ag ttt gc ag ttt gta ac ctt t.
```

147. An isolated promoter comprising the following nucleotide sequence:

ttggcaaactgagatataagaggggaaggtgattttcatgcaa
atTTTTTTTTTTTTTTTTgaatgaatgcaaaatttattcaaaaa
aaaaaacctgggctacatcaagtacttcatttctgagtttttgaaa
aatctaaagacaacaaaagactttacaatttaataaaaaaataataa
aaatactttatcactctcaacgaaattgttgatttaataacgtatct
cttggtaaaaacagcgttttatttgacgaaattgttataaatgaataa
aatgataatagaaaactagtgtggtacgtaaaaatacctctcatttggc
aaaataacggttatgtatcatgagatttgcatacgcacagcgtgctta
aatagtgtgctttcaggagaaaaatataccaagttatttgctgaaa
ttaccacgcaaactctgaggttcgaatggcaaaataaaaaaccaatgt
catttccttaatgtattaaggtcatttaataaaaattgtacactttt
ttcacctgtaagcgttccaaagtgtagaatggataactagaagggtc
aaaggtataatattaataagcgaactcactttttgcccaagtgattt
cacttcttacatttgcttgatatagttacccaaaagtgatatatat
tcccttatacaattgttctattttctggattataaggggaataagaa
aaaagaaaagagagagtatataataacttttataaagtgatgta
gatttctaatttgtaacgaaaagttcaaagtgaagaaaaaacgaaaa
agtttttctgttttgttttatatctatagccaagaaagtttctcaga
tttacaagaagttaactgagaaaaacaaaaaaaaaacttatgaagca
tgaaagactaattaacgaggtgatttaattttgagacaaattaaacat
cgaattaaaagtaacattttggaggggttatatgttatatatgtgaca
tgataagtcagattcatgactaatgtatatctggaatctaacatgga
agaatagagaacgaagcagagccaaggtcaacttgccagacacgaat
caacagatttgtgaatgagacaaatcaatgggtcataaacgggttggg
tttaaaccgggcaagtcaccttgggtcaattccattcggtatttcct
catgcaagaccctctgatacaaccaaaagactcccattacaatattct
ttcgatcacgagctacttattttcaaagtgtttacctctttcggtgac
tcttggtgtgtgtggttaaagcctagtcgagatgtgtcggtatatata
ggcatatatatacaaatgacgacaaaataagtatatttatattgtttaa
tttctatatattccatttctatatgcatgggtgggatttttgacaaaa
ccctaattcaagaatagaatccaaaagatgggatcaaagaatataat
ctaattgggctgaccacattttccgattttaattcgcatagttaatatt
ctttccactactttatgccgcagaaatttgtaattaagtaagacaaa
gaaatacagatatagatggtcgtagaaaccagtagaggaatttcat
tttctgtggataagtgggaatattaataagagaatggctcttactct
tacagtgggaaatgggaatagtagccattataaattcatcagattc
tatatatgcatgtttgtataagctaaaataaatacgtttaagcattc
ttcaaaaaaatttacaagttctagagactctcttaacgtcggcaatt
tatattctactttacatgacactttcaggaaaaagaaaactatactca
ctagcagatcattaaattttctttttctttttttgaatgaaccttag
ttgtgggtttttatttttggttagctagaaacttcagtggtttttttcc
gccaatggtagtgctttgatgatgggtccgg.

148. An isolated promoter comprising the following nucleotide sequence:

```
caatcaaggtaacgaaggaggatcagcgaaaggatgggcta
tatttggagttttttcctgcgtgtaagtaatgctttgtgatcttcca
tgcggacatataactgaagaataaactcaactcattgtgttctgggtg
tgtttcttctgatcagattcctcgttgcacatcgcacttttctgctgt
gggggctttatttataaaacaagagtagagcgtgtggtaatcttcat
atcttcttacaattccacttccattctcttaattattctctcacgtga
tatacacacactcaatcactgatgtactcgtatggatgcagcgtgga
actgatgcattggcggggatgtcacttctatcgggcttactagaaac
tgtaagtattacaagaaaactcaaaaggattccatttatgcaaaatc
taagagaaagctcactgtggtctttgggttacaatttatggatctctc
aagagacaaatgctatgtaagctaattgatttttgggtcttgataaaca
ggtagtggaagtggacaaagctactcaagaactgaagacatcaaca
atgcttttgccaatgaagtctcatgggaccgctcttccgcacatctct
actcaagcgacaacaacacagagaccaagtgaagaacatatgggtgc
gatctaattttgtcaagtgcctcacaagaggtagtgttcaagccat
ggtagggcacgcttgtgatctgcgatttctggattttgctttgtatg
tttattttctaccttctagaaagagggtcaaaaagttaatagcttcac
cgtgagaatgttgttttcaccagattcatgtgctatgatagaaaag
acaaagcaaacaagagttctttctttgcttaggttacaagaacaaga
gtatcgttataaagtcaacaaagattgaaacatatttttgtcaaggg
agtgggttagaatctcttctactctcttgcctttctcactaagacaa
aaaaaagacttggactttgtctaagggttttgtggatattattaacca
agtccttttgcaaaaagtaatatgttttttcgcattcctcttttag
aatttagtttaattctaggctttatatattgggttattactttcttgaaaa
atgatctgtttattctattcacttgggttacctcgctttttatctt
acttctacaaaaggattatcagtgaaggttagtctcttactctcacc
ttccgaaaataaaaacaaaatatcgatacttctagatcaaaccaagt
tgattaaaacatccctattccctaagattctgatcttgagatatatt
atcatgttaagatctaaattgacaagaaaactgatttttcatttcta
gtaggaaaaataattactatttagtgatcatgattgtcgaccgtaaga
ggtaggttagttactctccatctttctttgaagaagtcagaaagtca
gaaattatatcaaattaacatcaatattgaacacatatatctgtat
ggttttatgtttagaaaattccaatatatttatattcctagggaaaa
agaagcttattcttcaaattattgttatgagtcgttaaaatatggat
aaaaatataaagtctaaatatataaaaactcagttttgctttgctttta
cctctccaagctctccaaggtcaaatatttttagtttaattaaaccaa
aaaagggtttatttagtcaaacttagcatgcaatgctgggtaccaaacc
caagcatttagtctcttttaattcttctttttctccaataagtttttac
aatttttaattgtttgcatttcccttgattatttatcttcatcccaa
tttagctaataccaactccgtttcttattcttccaagtttttctta
taaatacgttcttcttccctcttatttcatatcactcaccacaaag
tcttctcatttctcat .
```

149. An isolated promoter comprising the following nucleotide sequence:

atgttgtgagtgaaggagaagaagagggaaacaaaggtatt
tatttgtagcgagttttgtttgtgacgcggtttgtctgtgttcaa
tgttgacgaaacgagtgagagagtgctgattattaaagaaaaccct
aattaagtcagacccgcccgttataaaaatagtcaaaaagtaggaaa
acgcgtgtgtgagtgtgacagagacagcccattgtttgctttatggg
cttataagcgagacgtgttaattgggctttttcctttatggccgaaa
acaaaagaaacgtgcctgagagattcgaactctcgccgggcagagcc
catgtacttagcaggcacacgccttaaccactcggccaaagcgactt
gttgctatgagtttagacaaaatcattaaaattctctattatgatttc
tcatagtgtgtgtgtatattgtggatctactaaaaattctttgttat
tattacttttattttgtgaattagtttgatataggttaagtacaaagtt
aactttattatttactcaaaaatttatcagattaactgattttatatt
gtttccttttggtatatatagacgtactatagtttttagaaaaaccataa
gattccttttatatttcatagagtgaagagatgagatgagatccttggc
tggagaagaaataagtttccacgaggaggactcttttttttttggtga
agacgaggaggaggactcttggttgatccagtctttacgttagacat
cgacccctacattttatttgcccttctctatcaacatggcaggtaaaa
atcttcattcaaccgaaccaaccaaaagtctcttcccaataatattca
agcaccatcctttgggaaactcatacatactacagtctacactcttt
cattttctttcaacgctcaacttaacaaatgatatagtctagttgtc
aattatatgttttaattagtgttttcacatcaaattctggtttgata
tttgatgactattttcggaaacatctcaatgtcccgc aaatacaatc
gtctatcatatataatcccgtacgttgtattctttatagatagaataa
tatggcgtgatctttataatataacatatagaatcgtgtagatttat
tttattttatttttatatatcgcataaattgcaaaatacttatatat
gtttgttatatatgatacccattttatagttacttaaaaaaagttaa
gcgataatatatatatatcaactttttataacaaaaaagtataacac
atggtaaaagaaaaataaaaatgaagacatgggtgtgacacgaaaatgg
cactaaatatacatatataatagatagctacaatatcccatcataca
cacttttttaattgactaatacataaacttacacacttttttaattga
ctaattcataactttttatcattgtcaacatgcaaattcatatttcc
gttgaactattattcttattttgtttttaaaagaagggtcttcctgggt
aataaaaaatgatgtttccaaatgacgttagagcaaaaaaaaaaaaaaag
gttgtctgggtctggtaaaatgaaaaagcaaagcgtcttgggtatagaa
aagtaatatactgcctcctaatttcttctgctccttctaccgaagaatc
tctccactcttgcctcttttcgaaaccctaaaccagaagcaccagat
tttttcaactttttccagagaacaatagaaaacccaacttgtgctc
tctagggtttttctttattccttctcatctttggattttcttgggtca
tcatttttggagcttaccaccagcgaaaaaattataacttccatcg
attcctgggtctctctctctcgctctctctgcatgtgctaaatcgccg
gactgatcctcactgtcacctctggt .

150. An isolated promoter comprising the following nucleotide sequence:

gattaggggtttgagttgtcactggaaagaggtttgattgt
gagtgatgatggagagattatgaaggagtttgtgtgtatttatagag
gagttaggggtttgaggtttgatgagaagtaggtttgaagaagtttt
gttgttgcaacttatttagagttacttgttccacaaccacaagtaag
attggtcacttctaagttctaactagaaacaaccatgacacatggag
atttcagctaacctagtttaatgtatatgtattatattttatttaaa
tattataaaaataaaaataaattttcacaaataaaaagaactacaaaaa
gtgagaaaaataatttgataaacaatttagaaaattagtatatcaa
taataaaattttataatccgatgggttttgcccttttggtttggcctttg
tttgaacttcgatgagtgactatgtatagcgaaaacaattcggtttg
tttttggtttaatttttaaaaaatacaagcgacaatatctgatgagaa
taggtgaaaagcaaataatatcagtttaattggaaatatctactttt
ttacaactaatattttgttttggtcaaccaacaatatagatttaattaa
ttatggtttatgagctttttatttgttgcgacagtatatatatgttaa
aatagtgatattgcatggcggaaggtccggaagcaacacatatctcc
tttttaatttttttttaacaagaataacatgttaatttttttttga
aattaataaagaatacatatttctaatttttgcgtcagatagatgat
taaagagtgtgtgttttttttaacaaacaaggaatacattatacata
tttcataatttctctcgacattgtttgttttttaaaaaatagattaa
agagtcctacgaagctaagtagctaacgaagacttgaaatgagaagaa
gacgagaatcttttaatatatttttgttaagcgataatattttga
ttaataaatatagattaaggaaataacaataacgcagatatcggtaa
gtcatagaaaaaaagaaacaacacaaacttacataaacatgtttcct
aatttglaatggagtaaaattccttcttttttttttttttttgattt
ggattccaattagtaaaagaactcaatgactataaataacctttaacc
ctctcattatttcttactatcaattgattaagctctcgttcctaaga
aagcaatagacgaacaagaacccatcgaagaacacaaatctctcttt
gaagttgtcgataatgttagtacaccgttacttcgtccaagactttt
ttgccgttcggtttcttacaaaacaaggatttggttaccattacttt
tgtcgtaactcctttttacatgtacgtcaaaaagtggttcctcgctc
cggccttgaagaaacgaccttcttaccacaaaaagcttatttttaaac
cgtctaaaaccggaaaatctcaatctaaaccggatacgggttcatgag
aaaccgattcaaacaccgagtgagaagtagaattttttgatgggtc
cgtcacaatgtgtgctgctccttcgccaagacatgtaccgattccga
tattttgtgggtgtaaagatgatcaaagagtccttcaaagctaagcacg
acttgaatgagaagaagaagaccaattactcaattagattttgtttt
gtggagcaattattgtctatttatctttgttttttagcaaataatctg
tatccactaatcttcacagtacttgactaacaagaagtaagaggttt
tcttattttccaattgttttttaatctgatacttttttcataatttta
caatgtttgatgaaaaaaacattcaaacctaaattttctttttttg
gtatgaattcaaacctgaattacttttgacgaggaccgacgggtata
aatagggtgatctcccaacaacaaaaagggt.

151. A transgenic plant or transgenic plant tissue comprising an isolated promoter according to any of claims 143 through 150.

54
APPENDIX 1

SEQ ID NO:	INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
1, 2, 3	2293133	glyceraldehyde-3-phosphate-dehydrogenase
4, 5, 6, 7	7143495	Histone H4
8 & 9	7143515	ATP dependent RNA helicase, mRNA sequence
10, 11, 12, 13	7143527	nematode specific
14 & 15	7143602	protein serine-threonine phosphatase 1, catalytic subunit
16 & 17	7143612	40S ribosomal protein S4
18	7143666	cytochrome p450
19, 20, 21, 22	7143675	Neuroendocrine protein 7B2
23, 24, 25	7143839	nematode specific
26	7143863	40S ribosomal protein S17
27 & 28	7144016	vacuolar ATP synthase subunit G
29	7144025	malate dehydrogenase
30 & 31	7144060	J2 pcDNAII Globodera rostochiensis cDNA similar to Bystin, mRNA sequence
32 & 33	7144225	similar to arginine kinase
34	7144354	pyrroline-5-carboxylate reductase

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
35, 36, 37, 38	C10	ribosomal protein L18a
39, 40, 41, 42, 43	C118	ribosomal protein S11
44 & 45	C122	ribosomal protein L16/L10E
46 & 47	C127	FMRFamide-related neuropeptide precursor
48	C129	ADP-ribosylation factor 1
49	C130	ribosomal protein L11
50	C137	nematode specific; conserved in <i>C.elegans</i>
51 & 52	C138	ribosomal protein L7
53	C145	ADP/ATP translocase
54 & 55	C148	troponin
56 & 57	C154	calponin
58	C16	translation elongation factor EF1A
59 & 60	C18	40S ribosomal protein S16
61	C27	ubiquitin
62 & 63	C46	nematode specific
64, 65, 66	C48	ribosomal protein S3AE
67	C59	40S ribosomal protein S5/S7

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
68	C8	glyceraldehyde 3-phosphate dehydrogenase
69 & 70	C82	60S ribosomal protein l30/L7E
71	C90	glyceraldehyde 3-phosphate dehydrogenase
72	C135	nematode specific
73 & 74	C206	predicted troponin
75	C227	cytochrome P450
76	C238	vacuolar ATP synthase subunit G
77	C246	40S ribosomal protein S4
78	C308	FMRFamide-like neuropeptide precursor
79	C342	ubiquitin
80 & 81	C344	nematode specific; conserved in <i>C.elegans</i>
82, 83, 84, 85	C370	40S ribosomal protein S5/S7
86	C426	nematode specific
87	C458	histone H4
88 & 89	C481	ribosomal protein L30E
90 & 91	C556	nematode specific; conserved in <i>C.elegans</i>

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
92	C628	ribosomal protein S17E
93 & 94	C665	malate dehydrogenase
95 & 96	C669	malate dehydrogenase
97	C694	ribosomal protein S3AE
98 & 99	C709	ADP/ATP translocase
100 & 101	C714	ADP-ribosylation factor 1
102	C721	calponin
103 & 104	C726	ribosomal protein L11
105	C736	nematode specific
106 & 107	C773	troponin
108	C834	nematode specific
109	C860	bystin
110 & 111	C863	troponin
112 & 113	C883	translation elongation factor eEF-1A
116	C888	40S ribosomal protein S16
117	C898	glyceraldehyde 3-phosphate dehydrogenase
118 & 119	C935	peptidyl-glycine alpha-amidating monooxygenase
120 & 121	C937	calponin
122 & 123	C942	peptidyl-glycine alpha-amidating monooxygenase

SEQ ID NO:	<u>APPENDIX 1 (cont.)</u> INTERNAL IDENTIFIER	FUNCTION OF POLYNUCLEOTID E / GENE
124	C954	arginine kinase
125, 126, 127	C969	calponin
128 & 129	7235653	ribosomal protein L18A
130	8005381	neuroendocrine protein
131	7235496	pyrroline-5-carboxyla te reductase
132 & 133	7275710	protein phosphatase pp1-beta catalytic subunit
134	7923685	nematode specific
135	7641370	40S ribosomal protein S11
136 & 137	7923404	nematode specific
138	7797811	ATP-dependent RNA helicase
139	7143613	predicted phospholipase D

Appendix 2:

Exemplary genes used for RNAi vectors.

Promoters:

Constitutive:

Super Ubiquitin from Pine

CCCGGGAAAACCCCT CACAAATACATA AAAAAAATTCTT TATTTAATTATC AAACCTCTCCACT ACCTT
TCC CACCAACCGTTA CAATCCTGAATG TTGGAAAAAACT AACTACATTGAT ATAAAAAACTA CATT
CTT CCTAAATCATAT CAAAATTGTATA AATATATCCACT CAAAGGAGTCTA GAAGATCCACTT GGACA
AATTGCCCATAGTTG GAAAGATGTTCA CCAAGTCAACAA GATTATCAATG GAAAAATCCATC TACCA
AACTTACTTTCAAGA AAATCCAAGGAT TATAGAGTAAAA AATCTATGTATT ATTAAGTCAAAA AGAAA
ACCAAAGTGAACAAA TATTGATGTACA AGTTTGAGAGGA TAAGACATTGGA ATCGTCTAACCA GGAGG
CGGAGGAATTCCCTA GACAGTTAAAAG TGGCCGGAATCC CGGTAAAAAAGA TTAAAATTTTTT TGTA
AGGGAGTGCTTGAAT CATGTTT'TTTAT GATGGAAATAGA TTCAGCACCATC AAAAAACATTGAG GACAC
CTAAAATTTTGAAGT TTAACAAAAATA AC'TTGGATCTAC AAAAAATCCGTAT CGGATTTTCTCT AAATA
TAACTAGAATTTTCA TAACTTTCAAAG CAACTCCTCCCC TAACCGTAAAAC TTTTCTACTTC ACCGT
TAATTACATTCCTTA AGAGTAGATAAA GAAATAAGTAA ATAAAAGTATTC ACAAACCAACAA TTTAT
TTCTTTTATTTACTT AAAAAACAAAA AGTTTATTTATT TTACTTAAATGG CATAATGACATA TCGGA
GAT CCCTCGAACGAG AATCTTTTATCT CCCTGGTTTGT AT'TAAAAAGTAA TTTATTGTGGGG TCCAC
GCGGAGTTGGAATCC TACAGACGCGCT TTACATACGTCT CGAGAAGCGTGA CGGATGTGCGAC CGGAT
GACCGTGATATAACCC ACCGACACAGCC AGCGCACAGTAT ACACGTGTCAAT TCTCTATTGGAA AATGT
CGTTGTTATCCCGC TGGTACGCAACC ACCGATGGTGAC AGGTCTGTCTGT GTCTGTGTCGCT AGCGG
GAGAAGGGTCTCATC CAACGCTATTAA ATACTCGCCTTC ACCGCGTTACTT CTCATCTTTTCT CTTGC
GTTGTATAATCAGTG CGATATTCTCAG AGAGCTTTTCAT TCAACCCGGG

Strawberry Banding Vein Virus 1

aagcttttcactgtgggttaatttcattaatctatccaggtgaaaacctcaaggaga
tctctcttctccaaaagacctctacagggcaatcaaaaactacagaaccagagttt
gtagtgcacagagtagaccaatctacctgagaatcacgagtaccttcttagagtggg
aaaatgatgacatccttattccataccactggattgaggtaggactatccaatggaa
aaattccatgggacaagtcatataagaagaccgcaacagtcgagtatcttcagaga
taactgcactcagacctaaaaggataaaagcagtatataatcagtgtagtaagatct
tcgcagattcaaagaagaagctt

Strawberry Banding Vein Virus 2

Gtttaaacacagcccaagataacagaaaagtc aaaggtgttcgaaagaccacttgt
gactaaggatcatttcatccataattatctggtagcacagactcatgataactgcga
ggaacacaagttctttacagtcgattcaaagacactttctctttacggtttcattga
aggagccgacccagaatatgtcagagaagcttttcactgtgggttaatttcattaat
ctatccaggtgaaaacctcaaggagatctctcttctccaaaagacctctacagggc
aatcaaaaactacagaaccagagttttagtgtagcagagtagaccaatctacctgag
aatcacgagtagcttctctagagtgaggaaaatgatgacatccttattccataccactg
gattgaggtaggactatccaatggaaaaattccatgggacaagtcatataagaagac
cgcaacagtcgagtatcttcagagataactgcactcagacctaaaaggataaaagc
agtatataatcagtgtagtaagatcttcgcagattcaaagaagaagcttaactatgc
tgatgacaagataattctaataagcaattattcagaattaatcaaggagaaagaatt
aataactctttcagaatatgaagcccgctttacaagtggccagctagctatcactga
aaagacagcaagacaatgggtgtctcgatgcaccagaaccacatctttgcagcagatg
tgaagcagcagagtggtccacaagacgcactcagaaaaggcatcttctaccgacac
agaaaaagacaaccacagctcatcatccaacatgtagactgtcgttatgcgtcggt
gaagataagactgaccccaggccagcactaaagaagaataatgcaagtggtcctag
ctccacttttagctttaataattatgtttcattattattctctgcttttgctctctat
ataaagagcttgatattttcatttgaaggcagaggcgaacacacacagaaacctccc
tgcttacaaacctgtattgttagctaaacctcttaggaggatatc

Nematode Inducible:**Trypsin Inhibitor from Arabidopsis (clone#6598343)**

ccccgggagcaaagcaagaacaccagagaagaagaaaagcactacagagaaaaatgtg
agcttaagcgcctctccaacaacacttctctgggagctctaaaggatgctgcaaaaagc
cttggtggtgagacttccgcataatctccaagcatgggtttatctttagcacaca
aactatctgacctcgacttggattttctctgcagtttgtccaactacattgaaac
ggatatgcaggcaacatgggatcatgaggtggccatctcgtaagattaacaaagtga
acaggtcactaaggaaaatacagacgggtactggactcgggtccaaggtgtagaaggag
gactaaagttcgactcagcaactggcgaattcattgcagttagaccttttattcaag
aaattgatacccaaaagggtctgtcgtctcttgataatgatgcacatgcaagaagaa
gtcaggaggatagcctgacgatacttcatcaagctccaggaagctaaatctgtcg
acaatgccattaagttagaggaggatataccatgaatcaagcaagaccaggtaga
acttctctatccataaaccatagatggagcgttagaatcttaattcattttcagtt
tttgaggatcattcatggagggttaattgctagtgtcagccatgggttggtggcc
aaagagctgtggcttgaaatggcagtggaagaaataaagagcgtttgcaacttaagctct
gtggaaatttcagatggaatggatccaacaatccgatgcagtggtcagttgttgaa
cctaaccaatccatgtcatgcagcatatcagattcatcaaattggctcaggcgcagtt
ctgcgtggaagctcatctacttccatggaagattggaaccaaatagagaaccacaaac
agtaatagcagcagagatggatcaacaacgctgatcgtaaggccagttatagagaa
gacactgtacgtttcaagttcgagccatcagttgggtgtcctcagctctacaaagaa
gttggaacacgttttaaaactgcaggacgggtcgtttcagctgaagtacttggtatgat
gaagaagaatgggtgatgctggttacagattctgatctccaagaatggtttggagata
ttacatgggtatgggaaaacactcgggtgaagtttctcgttcgtgattttgtctgccct
ctaggtagttctggtggcagtaattggttatcttggaacaggcttatgacgtcgtaag
acatagacacacacagttatgtattcccagtgaaagaatggtgtttatttctctaga
tattagtatgcttataaataggcatgaaggagaaagacaattttgggtatagtggagt
tcagcagaaaaatgtatatgtttttcgttttatatgaatcagagaataaaagtggga
tgttatatctacgtttgctaattgtgtacctgctcaccatctttcatataagaaaag
agaacacttttagttatccctgtgatgcagaatcgtaattctttgttatctctccatt
cctgtggaaaccaacaaagtcaactaaatttcgggttaattgggtgggttttaagtc
aacgaggacttgatttttagttgggtctgggcctataattgtgttcatttggtttt
tttcccccttatcagtttaacgtccatataccatctttttcttttttaacggcaa
agttcatatccatcttatgatgtgcctaaaagaggagagaagatgcgaagacagaa
ttttcatatttgaaagggttcgatatcgatattgggaaacgaatcaaggtcaaaaaa
ctcagtcataatagttgaaatttaaaaattttatttaattcaatccgattgggttcgtt
ttgttatgggttcggttctatatcatcaaaccaatcgggttggtcctaagataatta
taaatattcaccaacaccagtggttaaacacatatcaacaaacctaaagttagataaa
caaagagacccggg

**Arabidopsis Transmembrane Protein from Arabidopsis
(clone#6468048)**

ccccgggaattggcactcttctctgctgggttccaaaagaaacgaatcaatatgtgc
aacaagaagagctccagaagcagtcattctctaaaatcttaattctaacaacagctca
agaagaaaaaaattccatagctagagagaacacaaagtcaacagacgacgtcgtaga
ggcacaaagtcaaacctgaatggcttaagccgaactgagtggttttgactagaccat
catcagaaaaagctctccaagacggtagtcggatgttagatcgctcaagtaatttttgg
ttttgttggtctcacgttttcagctgccatttgatttcagtttggttttctctta
tctctaaaggcccaatttcatttaggttttagtttatttgatcattatccttactata
aaggctctcgctttcgagaaatttagggtttcttctgtctgtctcgctcactcaggtt
tgtgctcacaagactgcttcacttctagcttgattcttcttctcggtttatagtat
actgtacattagattattcttgtttctcgagcttctgctatagattttgattctttt
ttttgttgtctttgtttcgtttccaggatcagatcttagctaaattgagacaagctc
aaaatgaggtacttgacgcacatctctacattcactgtttaattagagaacaatacgt
ctctgaatcggtgattcagagacgtattgttcttctgtcatatgcaataagtttaatt
agagaacaatacgtctctgaatcggtgattgttttttggtatgtgcgttattgatagct

61

ttatgatgttaatagtctaggattgacacgaagtgttctgcagttttgcataaatg
ctctttactaaggcctctaaatttggatgacaaatctaaatcttgccctcataaaaaat
ttaggtgtattaagataagattatctttgtatggtagtgtctataatgtgggttgttc
atggttgaggttgtcaatgttgtgtatttttgtttgttttagttaatttgccttaactct
gttctttgtgggttaatacacagtaagcttcagagttagggcgttcgtgaagccatcac
tactatcacagggaaatccgaggcaaagaaacgtaactttgtcgagactattgagct
ccagatcggctctgaagaactatgacctcaaaaggacaagcgtttcagtggtatctgt
caagttaccacatatccccgtcctaaaatgaagatctgcatgctcggagatgccca
gcatgttgaagaggtgatatatcttttcatggaaattgatcattttgtgctctgttt
cttgtataatggttttgtgctcatttccatttgggtggctctatttagtttcattttagt
ttgtatctgtcttctgaatgtatagatgcatgattttcggaatttgggtcattgttta
tttaggcttcatttcttgcataattaaatatttgccttatttccatcttgtatcttttc
gtaggctgagaagatggggttggaaaacatggatgttgagtctctaaaaaagccttaa
caagaacaagaaactcgtcaagaagcttgcaagaaataccatgctttcttggcctc
tgagtctgtcattaagcagattcctcgtcttcttgggtcctgggtcttaacaaggcagg
caagttctggctacagctaataattccattgttcttctttacatccgttttgattttg
gataggtttttagtagtctatttcttttgtcaatgtctttttgatacaatgccaatcc
tttatcctgtgagattatgcttctttttagatgattcttaagtaacattcctttgttta
ctttacacaggaaaattcccaactcttgtgagccaccaggaatccttggagtc aaag
gtgaatgaaacaaaggcaacagtgaaagttccagctgaagaagggtctgtgcatggga
gttgcagttggtaacctttcccggt

**Diaminopimelate Decarboxylase from Arabidopsis
(clone#4159709)**

ccccgggtggcaaaactgagatataagaggggaagggtgattttcatgcaaatttttttt
tatttttttttgaatgaatgcaaaatttattcaaaaaaaaaaacctgggtacatc
aagtacttcatttctgagtttttgaaaaatctaaagacaacaaaagactttacaatt
taataaaaaaataataaaaaatactttatcactctcaacgaaattgttgatttaataa
cgtatctcttggtaaaacagcgtttttatgtgacgaaattgttataaatgaataaaat
gataatagaaaactagtgtggtacgtaaaaatacctctcatttggcaaaaataacggtta
tgtatcatgagatttgcatacagacagcgtgcttaaatagtgtgctttcaggagaaaa
tatataccaagttatttggctgaaattaccacgcaaatctgaggttcgaatggc caaa
taaaaaaaccaatgtcatttctttaatgtattaagggtcattttaaataaaaattgtacac
ttttttcacctgtaagcgttccaaagtgtagaatggataactagaagggtcaaagggt
ataatattaataagcgaactcactttttgcccgaagtgtttcacttcttacatttgc
ttgatatagttacc caaaagtgatatatatattcccttatacaattgttctattttct
ggattataaggggaataagaaaaaagaaaaagagagagtatataataactttttata
aagtgtatgttagatttctaatttgaacgaaaagttcaaagtgaagaaaaaacgaaa
aagtttttctgtttttgttttataatctatagccaagaaagtttctcagatttacaaga
agttaaactgagaaaaacaaaaaaacttatgaagcatgaaagactaattaacgag
gtgatttaattttgagacaaattaaacatcgaattaaaagtaacatttggagggttta
tatgttatatatgtgacatgataagtcgattcatgactaatgtatatctggaatct
aacatggaagaatagagaacgaagcagagccaagggtcaacttgcagacacgaatca
acagattgtgaatgagaccaaataatgggtcataaacgggttgggtttaaacgggca
agtcactccttgggtcaattccattcgttattccttcatgcaagaccctctgatacaa
ccaaagactcccattacaatatcttttcgatcacgagctacttattttcaaatgtgt
tacctctttcgtgactcttgtgttgtgtggttaaagcctagtgcgagatgtgtcggtat
atataggcatacatatacaaatgcgacaaaataagtatatttatattgtttaatttct
atattccatttctatatgcatgggtgggtattttgaccaaaccctaattcaagaat
agaatccaaaagatgggtcaaagaatataatctaattgggtgaccacattttccga
tttaattcgcatagttaatatctttccactactttatgccgcagaaatttgaatt
aagtaagacaaagaaatacagatataagatgggtcgtagaaccagtagaggaatttc
attttcgtggataagtggaatattaataagagaatgggtctttactctttacagtgg
gaaatgggatagtagcccattataatttcatcagattctatatatgcatgtttgta
taagctaaaaataaatacgttttaagcattcttcaaaaaaattttacaagttctagagac
tctcttaacgtcggcaattttataattctactttacatgacactttcaggaaaagaaaa
ctatactcactagcagatcattaaattttctttttcttttttttgaatgaaccttagt
tgtgggtttttatttttgttagctagaaacttcagtggtttttttttccgccaatggtag

tgctttgatgatggtccggcccg

Peroxidase from Arabidopsis (clone#4006885)

cccggaatcaaggaacgaaggaggatcagcgaaggatgggctatatattggagt
tttttcctgctgtaagtaaatgctttgtgatcttccatgaggacatataactgaaga
ataaactcaactcattgtgttctgggtgtgtttcttctgatcagattcctcgttgcat
ctgcacttttctgctgtgggggctttatttataaaacaagagtagagcgtgtggttaa
tcttcataatctttctacaattccacttccattctctaattattctctcacgtgatat
acacacactcaatcactgatgtactcgtatggatgcagcgtggaactgatgcattgc
cggggatgtcacttctatcgggcttactagaaaactgtaagtattacaagaaaactca
aaaggattccatttctgcaaaatctaagagaaagctcactgtggcttttgggttaca
tttttggtatctctcaagagacaaatgctatgtaagctaattgatttttgggtcttgata
aacaggtgagtgggaagtggacaaagctactcaagaactgaagacatcaacaatgctt
ttgccaatgaagtctcatgggaccgctcttccgcatcttctactcaagcgacaacaa
cacagagaccaagtgaagaacatatgggtgcgatctaattttgtcaagtgccctaca
agaggtactgtttcaagccatgggtatggcacgcttgtgatctgcgatttctggattt
tgctttgtatgtttattttctaccttctagaaaagggtcaaaaagttaatagcttca
ccgtgagaatgttgttttaccagattcatgtgctatgatagaaaagacaaagcaa
acaagagtcttttcttggcttaggtttacagaacaagagtatcggtataaaagctaac
aaagattgaacatatattttgtcaagggagtggttagaatctcttctactctcttg
cctttctcactaagacaaaaaaagacttggactttgtctaagggttttgtggatatt
attaaccaagtccttttgcaaaaagtaatatgttttttgcattcctcttttagaa
tttagtttaatctaggctttatattgggttattactttcttgaaaaatgatctgttta
ttctattcatacttgggttacctcgtttttatcttacttctacaaaaggattatcag
tgaaagttagtctcttactctcaccttccgaaaaataaaacaaaaatcgcatacttc
tagatcaaaccaagttgattaaaacatccctattccctacgattctgatcttgagat
atattatcatgttaagatctaatttgacaagaaaactgatttttcatttctagtagg
aaaaataattactatttagtgatcatgattgtcgaccgtaagaggtgggttttagttact
ctccatctttctttgaagaagtcaaaaagtcagaaattatatcaaatcaaacatcaa
tattgaacacatatatctgtatgggtttatgtttagaaaaattccaatatttatatat
tcctagggaaaaaagaagcttattcttcaaattattgttatgagtcgttaaaatattg
ataaaaaataaagtcataatattaaaaaactcagtttgcctttgcttttacctctca
agctccaaagtcataatatttttagtttaattaaacaaaaaagggttttagtcaaa
acttagcatgcaatgctgggtaccaaaccgaagcattagtcctcttttaattctcttt
ttctccaataagtttttacaatttttaattgtttgcatttcccttgattatttatct
tcatcccaatttagctaataccaactccggttcttattcttccaagtccttttccat
aaatacgttcttcttccctcttattttcatatcactcaccacaaagtccttctcattt
cctcatcccg

**Mitochondrial Uncoupler from Arabidopsis
(clone#4220510)**

cccggaatggtgtgagtgaggagaagaagagggaaacaaaggtattttattttagtc
gagttttgtttgtgacgcggtttttgtctgtgttcaatgttgacgaaacgagtgaga
gagtgctgattattaaagaaaaccctaattaagtacagaccgcggttataaaaaat
agtcaaaaagtaggaaaacgcgtgtgtgagtgagacagagacagccattgtttgct
ttatgggcttataagcgagacgtgttaattgggctttttctttatggccgaaaaca
aaagaaacgtcgctgagagattcgaactctcgcgggcagagcccatgtacttagca
ggcacacgccttaaccactcgccaaagcgacttgttgcctatgagttagacaaaatc
attaaaattctctattatgatttctcatagtggtgtgttatattgtggatctactaa
aaattctttgttattattactttattttgtgaattagtttgatataggtaagtacaa
agttaaactttatttttactcaaaatttatcagattaaactgattttatattgtttcc
tttgggtatatagacgtactatagtttttagaaaaaccataagattccttttatattc
atagagtgaagagatgagatgagatcttggctggagaagaaataagtttccacgagg
aggactcttttttttgggtgaagacgaggaggagactcttgggtgatccagctctt
acgttagacatcgacccctacattttatttgcctttctctatcaacatggcaggtaaa
aatcttcattcaaccgaaccaacaaagtccttcccaataatattcaagcaccatc
ctttgggaaactcatatactacagtcctacactctttcattttctttcaacgctca
acttaacaaatgatatagtctagttgtcaattatatgttttaattagtgttttcaca

63

tcaaattctggtttgatatttgatgactatcttcggaacatctcaatgtcccgc
atacaatcgtctatcatatataatcccgtacgttgattcttatagatagaataa
tgccgtgatctttataatataacatatagaatcgtgtagattttattttatttt
ttatatatcgcataaattgcataaatacttatatatgtttgttatatatgataccat
tttatagttacttaaaaaaagttaagcgataatatatatatcaactttttataac
aaaaaagtataacacatggtaaaagaaaaataaaaaatgaagacatgggtgtgacacgaa
aatggcactaaatatacatatataatagatagctacaatatcccatcatcacactt
ttttaattgactaatacataacttacacacttttttaattgactaattcataacttt
ttatcattgtcaacatgcaaattcatatttccgttgaaactattattcttattttgtt
tttaaaaagaagggtcttccgtgtaataaaaaatgatgttccaaatgacggttagagcaa
aaaaaaaagaagggttctggtctggttaaaatgaaaaagcaaacggtcttggtatag
aaaagtaataatactgcctcctaatttcttctgctcttaccgaagaatctctccact
cttgccctcttctcgaaaccctaaaccagaagcaccagattttttcaactttttccca
gagaacaatagaaaacccaacttgtgctctctaggggttttctttattcttctctc
tttggattttcttgggtcatcattttgggaagcttaccaccagcgaaaaaattataa
cttccatcgattcctgggtctctctctcgcctctctctgcatgtgctaaatcgccgg
actgatcctcactgtcacctctgttcccggg

Stress protein from Arabidopsis (clone#6598614)

ccccggggattaggggtttgagttgtcactggaaagagggtttgattgtgagtgatgat
ggagagattatgaaggagtttgtgtgtattttatagaggagttagggttttgaggttt
gatgagaagtaggtttgaagaagttttgttgttgcaacttatttagagttacttggt
ccacaaccacaagtaagattgggtcacttctaagttctaactagaaacaacctatgaca
catggagattttcagctaacctagtttaattgtatatgtattatattttatttaaatat
tataaaataaaacataatttccaaaataaaagaactcaaaaaaagtgaaaaaataa
tttgataaaacaaatttagaaaaattagtatatcaataaataaatttataaatccgatgg
ttttgccttttgggtttggcctttgtttgaacttcgatgagtgactatgtatagcgaa
aacaattcgggtttgtttttgggttaattttaaaaaatacaagcgacaatatctgatg
agaataggtgaaaagcaataatcagtttaattggaaatatttacttttttaca
ctaataattttgtttgggtcaaccaacaaatagatttaatttaattatgggtttatgagct
tttatttggttgcgacagtatatatatgtttaaatagtgatattgcatggcggaagggt
ccggaagcaacacatatctccttttttaatttttttttttaacaagaataacatgttaa
tttttttttgaatttaataaagaatacatatttctaatttttgcgtcagatagatga
ttaagagtggtgtgttttttttaacaacaaggaatacattatacatatttcatatt
tctctcgacattgtttgttttttttaaaaaatagattaaagagttctacgaagctaagt
agctaacgaagacttgaaatgagaagaagacgagaatcttttaatatatttttgttaa
gcgataatatatttgaaaattataaatatagattaaaggaaataacaataacgcagat
atcggttaagtcatagaaaaaaagaacaacacaaacttacataaacatgtttcctaa
tttgaataaggagtaaaattccttcttttttttttttttttggatttgattccaatta
gtaaagaactcaatgactataaataacctttaaccctctcattatttcttactatca
attgattaagctctcgttcctaagaagcaatagacgaacaagaacccatcgaagaa
cacaattctctcttgaagttgtcgataatgttagtacaccgttacttcgtccaaga
ctttttttgcggttcggtttcttcaaaaacaaggatttggttaccattacttttgcg
taactcctttttacatgtacgtcaaaaagtgggtcctcgcctcgggttgaagaaacg
accttcttaccacaaaaagcttatttttaaccgtctaaaaccggaaaatctcaatc
taaaccggatacgggttcagagaaccgattcaaacaccgagtgagaagtagaatt
ttttgatgggttcggtcacaatgtgtgctgctccttcgccaagacatgtaccgattcc
gatattttgtgggtgtaagatgatcaaagagttctcaaagctaagcagcacttgaat
gagaagaagaagaccaattactcaattagattttgttttgggagcaattattgtct
atttatctttgttttttagcaataatctgtatccactaatcttcacagtacttgact
aacaagaagtaagagttttcttattttccaattgttttttaattctgatacttttttc
ataattttacaatgtttgatgaaaaaaaacattcaaacctaaattttcttttttgg
tatgaattcaaacctgaattacttttgacgaggaccgacgggtataaatagggtgat
ctcccaacaaacaaaaagggtcccggg

Pectinacetylsterase from Arabidopsis

(clone#6671954)

ccccgggtgggtggggacaatggatccggtctgcgtagcaacaaggctgaaaaagatta




FIG. 1




FIG. 2




FIG. 3




FIG. 4

Selectable Markers

pNOS / NPT-II / tNOS

pSU / Bar / tNOS

pSU/ Intron / Bar / tNOS

pUBQ3 / Intron / PMI / tNOS





FIG. 5

Scorable Markers



¹ Construct useful for promoter analysis.

² Construct useful for high constitutive expression of genes of interest.

FIG. 6





FIG. 7



* RNAi vector adapted from MBCS Shuttle vectors
pAKK1413 and/or pAKK1414 (Figure 2).

FIG. 8

AKK110P1
SEQUENCE LISTING

<110> Mushegian, Arcady R.
Taylor, Christopher G.
Feitelson, Gerald S.
Eroshkin, Alexey M.

<120> Materials and Methods for RNAi Control of Nematodes

<130> AKK-110P

<140>

<141>

<160> 139

<170> PatentIn Ver. 2.1

<210> 1

<211> 165

<212> DNA

<213> Globodera rostochiensis

<400> 1

```
gtttgagatt attgactttg catattttcca accaagttca tttagaccaat attttcctgc 60
taaacaatagc aaaaatggtg aaaccgaagg tcggcattaa tggcttttga cgcattgggc 120
gcttggcggtt gcgcgctgcg gttgagaagg acaccgttca ggtgg 165
```

<210> 2

<211> 342

<212> DNA

<213> Globodera rostochiensis

<400> 2

```
cgactacatg gtatacatgt tcaactacga ctcgacccat ggccgcttca atggcaaaaat 60
ttcgacaagc gccggcaatt tggtcgttga gaaagagggg aaggccacgc acaccatcaa 120
ggtgttcaac ctcaaggacc cggccgagat caaatgggct gaggtgggcg cggaatatgt 180
gatcgagtcc accgggggtgt tcaactaccat tgagaaggct tcggcacact tgaagggggg 240
cgccaagaag gtggtcatct ctgctccgtc cgctgatgca ccgatgtacg tgatgggctg 300
caacgaggac aaatatgacc cggccaagga caacgtgatt ag 342
```

<210> 3

<211> 205

<212> DNA

<213> Globodera rostochiensis

<400> 3

```
gaagccggcc tcattggacg ccatcaaggc ggcggtgaag aaggctgccg aagggaattt 60
gaagggcatt ttgggttaca cagaggacca ggtggtgtcc acggactttc ttggagacag 120
tcgctcgtcg atcttcgacg ctggggcggtg catctcgttg aaccgcact ttgtcaagtt 180
ggtcagctgg tacgacaatg aattt 205
```

<210> 4

<211> 167

<212> DNA

<213> Globodera rostochiensis

<400> 4

```
ttaaagcatt tattcacacg cacggagaaa tgaggattac ctaatttgat tgagtctttc 60
tcgtccattt gtcaattgtg gccctaaaga gggccggttg ggtagttttt ttggtgttcc 120
ttctccttgc tggctcaacc accgaagccg tacagcgtcc ggccttg 167
```

<210> 5

AKK110P1

<211> 41
 <212> DNA
 <213> Globodera rostochiensis

<400> 5
 catggccgtc acggtcttgc gcttcgcgtg ttcgcagtat g 41

<210> 6
 <211> 79
 <212> DNA
 <213> Globodera rostochiensis

<400> 6
 gtttcccagg aaaactttca gcacggaacg agtctcctcg taaatgaggc cagagatgcg 60
 cttaacgcct ccacgacgg 79

<210> 7
 <211> 168
 <212> DNA
 <213> Globodera rostochiensis

<400> 7
 cggcttggtg atgccctgga tggtatcccg caagactttt cggtggcgtt ttgcgcctcc 60
 ctttccgagt ctttttccgc ctttcccgcg tccggacatt ttgttggtta atcagaagag 120
 cacagagagt aggagaaata ggaaattttg cctcgtgccg aacgtgcc 168

<210> 8
 <211> 330
 <212> DNA
 <213> Globodera rostochiensis

<400> 8
 gacagtctcc gttctgggta tgtgtcacac gcgcgaactt gctttccaaa tttctaagga 60
 atacgagcga ttcaccaagt acatgccggg agtgaagggt tccgtattct tcggagggat 120
 gccgataaag aaagacgaag aggtattggc taagaacacg ccgcacattg tcgtcggaac 180
 gccgggacgt cttttggcct taggacgcac tggacatctg aagctgaaag gcgtcaaata 240
 ctttgtgctg gacgaatgcg acaaaatgat tggagatgcc gacatgcgcc acgacgtgca 300
 ggaaatcttc aaaatgacgc ctcaggagaa 330

<210> 9
 <211> 136
 <212> DNA
 <213> Globodera rostochiensis

<400> 9
 actttgccgc gggagctgcg cgtcttctgc aaaaagttca tgcaggaacc aatggaggta 60
 tacgtcgacg acgaggctaa gcttacgctt cacggtctcc aacaatacta cgttagactg 120
 aaggaaaatg agaaga 136

<210> 10
 <211> 141
 <212> DNA
 <213> Globodera rostochiensis

<400> 10
 tattaaaata aaatacaaac aataatataa tggctgtttt ttctgtcatg tttaagttt 60
 ttgttggtca tcaactttct cagcagcgac aatacggcca atccggtgaa agggccaaag 120
 tcaatagctc gtcggtacc t 141

<210> 11
 <211> 141
 <212> DNA

AKK110P1

<213> Globodera rostochiensis

<400> 11

```

aaccaggcac tctgttcac ttcggcatcg ctttttggca atgtcaacaa cactttgctg 60
gccattttgt ttctacagca cacgcacacc gtcgtcttta cagcgttcac ctcgcaaaa 120
aagtagccgt atttgcgaaa t 141

```

<210> 12

<211> 37

<212> DNA

<213> Globodera rostochiensis

<400> 12

```

gcgttggtg caagctgtac acaaggctgc ccggttt 37

```

<210> 13

<211> 161

<212> DNA

<213> Globodera rostochiensis

<400> 13

```

gcgcgttcca tcgcccgcac cacaaaaagt cccatcgctt catatcgtag cgcaaatgt 60
ctttggtgca aatggcaaaa cggccaaaat aatggtcgaa gccgtacaca accgccaccg 120
ccacagcgcc aacccacac caaatgcgaa atttatcgaa a 161

```

<210> 14

<211> 306

<212> DNA

<213> Globodera rostochiensis

<400> 14

```

gaattcgttt gaggtataaa taaataataa atggcagcca acgaatcgct aaatgtggac 60
agtttgatca ctcgattgtt agaagttcgg ggttgtagac cgggaaaaac agtgcaaatg 120
gacgaatctg agatacgcac tttgtgcac aaaacacgtg aaattttgct gtcgcagcca 180
atcttggttg agctcgaggc accittaaaa atttgtggtg acattcacgg acaatataat 240
gatcttctga gattgttcga atatggtggg tttccaccgg aagcgaacta tctatttctt 300
ggggac 306

```

<210> 15

<211> 261

<212> DNA

<213> Globodera rostochiensis

<400> 15

```

gcaaagcctt gagacgattt gtttgcctgt tgcttacaag attaaatatt ctgaaaattt 60
ttttcttctt cgtggcaatc acgaatgtgc ttcaatcaat cggatttacg gattttatga 120
tgaatgcaaa cggagggttc tcaatcaagt tgtggaagac cttcactgac tgcttcaact 180
gtctgccaat tgccgcttta atcgacgaaa agatcttttg ctgccacgga ggctgtctcc 240
tgatttgcta aacatggcag c 261

```

<210> 16

<211> 151

<212> DNA

<213> Globodera rostochiensis

<400> 16

```

gaattctttg agtgcattca gcgtttaatt ttttcgtatt ataataagca tggctcgcgg 60
acccaaaaag catttgaagc gacttgcagc acccaaaaaa tggatgttgg acaaattggg 120
tggcgttttt gcgccacgtc cattgtgcgg a 151

```

<210> 17

<211> 306

AKK110P1

<212> DNA

<213> *Globodera rostochiensis*

<400> 17

tcaagtacgc	gcagtcgtac	aacgaagcgc	gcatgatctg	caaacagcgg	ctgatcaagg	60
tggacggcaa	agtgcgcacc	gagatgcgct	tcccgtgcgg	aataatggat	gtgatctcga	120
ttgagaagac	aaacgaaacg	tttcgtctgg	tgtacgatgt	gaagggccgt	tttgtcatcc	180
atcgaattca	aaagctggag	ggccagtaca	agctgtgcaa	agtgaagaag	caggccgtcg	240
gggacaagca	ggtccccctac	attgtcacac	atgacgcgcg	caccattcgc	taccggaccg	300
ctcatc						306

<210> 18

<211> 528

<212> DNA

<213> *Globodera rostochiensis*

<400> 18

gaattcgcac	aacgaattga	agacttatgc	ggcagaaaaa	ggacttttgc	caaagtgtga	60
ggagcaagca	gacgaccttt	cggattggct	ttgttcgtcc	attgggttgg	agcatcggcc	120
gttcctaccg	tatacaaacg	ctgtaataaa	tgaacaatt	cgattagtca	at ttgatccc	180
gttcaatctt	agccatttgg	cgcttgaaga	tatgcaaat	ggcaatttta	ttgtgaagcg	240
tgggacacca	attgtaccgc	aggtcagcag	tgttctgttc	gacgaaaaac	tgtatccgga	300
gcccgatcgg	tttttgcccg	aacgctttct	ggacgatgag	ggccgtttga	agaaaagcga	360
cgaacttatt	gcatttgggg	ttgggaaaag	gcaatgtgcc	ggcgaagctt	tggcccgaat	420
gacacttttt	ctgtttgccc	ctaatttctt	tctcgcctac	aaagttctcc	cgctccgatcc	480
actgaatcct	ccaagcctga	aaaagttggc	ggattatctg	tttacaca		528

<210> 19

<211> 335

<212> DNA

<213> *Globodera rostochiensis*

<400> 19

gaattctttt	agaaagcggg	aattcgtttt	tggctataaa	atgattctgt	gggccacgat	60
tttgttgatg	gctttggaca	ttgcgttcgg	tggcaccaat	caaattggaat	ttgatcagtc	120
ggcgccgatg	ttccccgact	cccagttcat	cgatttgatt	tcgcgcgaca	tcgaatcctt	180
ctccggccca	ttgggcgttg	gccataaatt	tatgagcggc	ggtgccggtg	agggcgctcca	240
acagctaggc	cccgaagggc	cctttgagca	gcggcaacag	gtgaagagtg	acaatgttct	300
ccccgcgtat	tgcgagcctc	caaatccctg	tccga			335

<210> 20

<211> 52

<212> DNA

<213> *Globodera rostochiensis*

<400> 20

ggacggctgc	acggaacagt	tcgagaacac	tgccgagttt	tcgcgcagct	ac	52
------------	------------	------------	------------	------------	----	----

<210> 21

<211> 190

<212> DNA

<213> *Globodera rostochiensis*

<400> 21

gcttgtgtga	ccaggagcac	atgttttaact	gtccgtcgaa	gaacaaccgc	gaggagtacg	60
agcaggatct	ggagcaattg	ctggccaaca	acggactgca	caaataaatg	attgccaaga	120
aattccatct	cacgcggggc	gaggagccgc	gccgtcgaaa	acgctcttgt	cgcccggctt	180
cggccaaccg						190

<210> 22

<211> 52

<212> DNA

<213> *Globodera rostochiensis*

AKK110P1

<400> 22
ccgctacaac ccctacctgg agggcgcccc gctgaagtca gtggccaaaa ag 52

<210> 23
<211> 54
<212> DNA
<213> Globodera rostochiensis

<400> 23
gaattccgac tctcaaggtg gacccacgcc caaccaacag caattgtcag ctgc 54

<210> 24
<211> 77
<212> DNA
<213> Globodera rostochiensis

<400> 24
ccgcacatgt cgaggcctcc atcttttggc actgggtcatc accttccgcc tactgctaac 60
aacagaccgg aacagca 77

<210> 25
<211> 439
<212> DNA
<213> Globodera rostochiensis

<400> 25
gtcaatcaaa aacgccgact tgcattcctc agctgatggt cagtaatgcg ctaaagggca 60
tccattccgt ctcttctaca tcagcaacac aatcacattc cagccccagt tttatgacac 120
acaacgtgca gcagcaacat gttgttggtc aacaacagca gcaacaacag aatttccaac 180
aaccgccggc cctatcgtag actcacagcc accaacaaca aaaacaacca ccacaagcgt 240
cacagtcgat gttgtcaatg aaaagtggca atgttggtcgt tgttggtccg caacaatcgc 300
agcagcacca ctaccaacag cggacactga cgccactgaa gcacacatcc gcacacctca 360
cgtccgatcg cttcgtcatc accaaaacca acagggtgct tccactcccg tcgcagcaag 420
gcgccacggc cactgatga 439

<210> 26
<211> 539
<212> DNA
<213> Globodera rostochiensis

<400> 26
gaattcggtt gagacacatc caattaatta atagttattg ttggcaatgg gacgagttcg 60
aacgaaaact gtgaagaagg cgtcgcgcgt cattattgag aagtattaca ccaaattggg 120
cctcgacttt cacaccaaca agcgcatttg cgaggagggt gccattatcc caagcaaacg 180
gatgcggaac cgaattgcgg gatttatcac acatctgatg aagcgcattg agctgggccc 240
tgtccgtggc atttccatca aattgcagga ggaggagcgc gagcgtcgcg acaattacat 300
gcccgaatc tcttacctgg atgcgcagaa tcaccagatg atcagcaccg accaagagac 360
gaaggatatg gcggaatttc tggggctagg cctcaacttg gaagtgaag ggcctttgac 420
gagtggcggc gctggcgtag gacgtcgttg agtcaggaca attggcatta ttgttgaaaa 480
atcatcgatg tttgttcgc atttgatga taatgcgctg ataaattttt gttgatttt 539

<210> 27
<211> 179
<212> DNA
<213> Globodera rostochiensis

<400> 27
gaattcnaca gtttctgtga gtaatggcat ntcacactgc cggcatccaa cagttgcttg 60
cggccgaaaa gcgtgcggca gaaaagatta atgatgccc gaagcgaaaa gcacagcgac 120
ttaagcaggc caaacaagaa gccaggcgag agatcgagca gtatcgncag gagagggag 179

AKK110P1

<210> 28
 <211> 133
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 28
 gcaaaattat ttgggcacgc gcgacgacat cgagcagcaa ataaagcgcg agacagaaga 60
 gtcgctggag gcaatgaatc gcaatgtcgc ggcgaacaaa cagcagggtca ttgtacgtct 120
 gctgcagttg gtg 133

<210> 29
 <211> 482
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 29
 gaattcgtga aatcaaaagc ttttttaatt tatttacaca aaaaatgggt ccaccaccaa 60
 ttgcgctggt ggtcactggt gccgctggac aaattggcta ttcactgggt ctgcaaatcg 120
 caaaaggcga tgtgtttggc aaagatcagc caattgttct cgttctctc gacattccac 180
 cgatggccga agtactctct ggtgtccatt ttgaattgat ggactgtgctg ttggcaaac 240
 ttgccgggtgt ggaggctgtg accacggaag agcaggcctt caaggacatt gactacgctt 300
 ttctttgtcgg agcgatgccc cgaagagagg gaattggaacg aaaggacctt ttggcggcaa 360
 atgtcaaaat tttcaagtcc caaggcgaag cattggcccg cttttccaag cccgtncgtc 420
 aaagtctctg tgggtgggcaa cccggccaac acgaacgcgt acatttgctc aaaatatgcc 480
 gg 482

<210> 30
 <211> 605
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 30
 gaattcaaag tgccgaaagc gttcaaaata attccggcaa tggcaaattg ggaacaaatt 60
 ctagacctta cttccccga aaaatggagt tcagcggcga tgtttcaagc aactcgtgtg 120
 ttttctgccca cgggcacacc gtcacaatgc caaagggtta acactttggt gctgttgcca 180
 cgactccgtg atgagattga cgagtacaag aagctaaact ttcatttgta tcagtgtgtg 240
 tttaaagcaa tgttcaagcc ggccggattt ttttaaggga ttattttgcc tctttgcaa 300
 tctggcactt gcaactctcg tgaagccatc atctttgggt ctgctctgctg aaagatttca 360
 ataccgcaac tccacgccgc tgcagcaatg ctcagcatag caaaaatgga ctactcgggc 420
 gccatttctt ttatcctacg tgttcttggg gaaaaaaatt acacacttcc tttccgagca 480
 tttagcggcc tcgtttttca ttttcttggg atgcgctcac atcagggcga gctgccagt 540
 atttggcacc agacactgtt ggcttttgtc gagcggttac caaaagacat aagtgcagaa 600
 cagag 605

<210> 31
 <211> 112
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 31
 ccattcccat catcaaatta ccccgattta ctgcggcttt tgcgcggcgc cgagtcgagg 60
 aatgaggaat gtgaagcaaa tgtgcccgtt tatgcgcgta atgatgaaat gg 112

<210> 32
 <211> 105
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 32
 gaattcgttt gagcatttat ttgacaaaat ctgaataaat ggccgtacca aaagaagtta 60
 ttgacaaaat cgaggcgggt tacaagaagc ttcagggaagc gtctn 105

<210> 33

AKK110P1

<211> 425
 <212> DNA
 <213> Globodera rostochiensis

<400> 33
 aagaagtacc tcaccaagga agtcgtcgat gcctgcaagg ataagcgac caagcttggg 60
 gcgaccttg cggatgtgat ccagtcgggc gttgccact tggacagcgg agttgggggtg 120
 tacgctcctg acgctgaggg ttacaccttg ttcaagccgt tggtcgaccc gatcatcaac 180
 gactaccatg gtggcttttg tccgggcagc aagcagccgg caactgacct tggtagacggc 240
 aaaacgcana tgctgaccgg atctcgaccc cgaggggaaa atttatcaat ttcgacacgc 300
 gttcggttgcg gccgtttcct ttaagggata cccggttcaa cccgtgcttg acnaaaggan 360
 aactacnttt ggagatggga aacnaaggtc nagggccgtt ttctaacatt ttnaagggn 420
 atcct 425

<210> 34
 <211> 581
 <212> DNA
 <213> Globodera rostochiensis

<400> 34
 gaattcggtt gagcgaagag ttttgtggtt gacaccgggt tatggacttt tagcccggtga 60
 tccttgacgg tccaaagccg cgttcagttc cgtgccgtgt tttttaaaag aggcgggagag 120
 tttgacggtc attccaagca gccaaataaac caccaaaacc aaataacccc cccaatcga 180
 tccccccct ccaattcctc cgcattattc gcattatcaa ttctaatacag cacaaccact 240
 gcatcattcc tttcccgaac atacgatgct aagtgaact ttgaaaattg gcttcacg 300
 agccggaaag atggcccaag cattggcaag aggacttatt aattcggggc gatacccggc 360
 agagaatttg atggcgagtt gtccaaagac ggacgaggct ttactggagc aatgcaaaaa 420
 attgggaatc ggaacgacgc acgacaacac tttggtcgcg cgagagaacg acgtcatcgt 480
 attggcggtc aagccgatgc acatcagcaa agtgacgtcg gaaatcgac ccaatttccg 540
 gaggaacat ttgcttattt cattgattag gaattacact t 581

<210> 35
 <211> 102
 <212> DNA
 <213> Globodera rostochiensis

<400> 35
 gaattcggtt gagaatttta ctttatataa ttgacgttta atcagcagcc ataagcaatg 60
 cccatcaaag catccggaga aacattaagg aagtttattg tc 102

<210> 36
 <211> 34
 <212> DNA
 <213> Globodera rostochiensis

<400> 36
 tgcaaagat gcaaaccaca cgcttcacaa gatg 34

<210> 37
 <211> 100
 <212> DNA
 <213> Globodera rostochiensis

<400> 37
 tcatgttggt gccaaatctc gcttctggta ctttacgagc atgctgcgtc gagttaagaa 60
 aacacacgga gagatcggtt cgtgtcaaga ggttttcgag 100

<210> 38
 <211> 176
 <212> DNA
 <213> Globodera rostochiensis

<400> 38

AKK110P1

tgaagaactt cggaatttgg ctccgttacg attctcgtac tggacaccac aatatgtacc 60
 gcgagtatcg ctgatgttac cgaggccggt gccgtgacct aatgctatcg cgacatgggc 120
 gctcgtcacc gcgctcaggc ggatcgaatt caaatcatca aagtgcaaac ctcaag 176

<210> 39
 <211> 155
 <212> DNA
 <213> Globodera rostochiensis

<400> 39
 gaattccaag tttgaggtat tgtttgttat acgatttctt acaaatgaca gaacaaactg 60
 agcgcgcgtt ccaaaaacaa ccgatcggtt ttctgaacga caagttcaga acgcaaggga 120
 ttgggaagaa ggcattccaac aaggaccgtt actgg 155

<210> 40
 <211> 35
 <212> DNA
 <213> Globodera rostochiensis

<400> 40
 tcctcgcgag gctattgagg gcatatatat cgaca 35

<210> 41
 <211> 70
 <212> DNA
 <213> Globodera rostochiensis

<400> 41
 tggaaatgtg tccatccgcg gtcgcattct cactgggggtg gtgatcaaaa acaaaatgca 60
 gcggacgatt 70

<210> 42
 <211> 85
 <212> DNA
 <213> Globodera rostochiensis

<400> 42
 tcgtaccaaa atatcgtcgc tatgagaaac gccacaaaaa catgtccgtc cactgttcgc 60
 cgtgcttcgc agatgtctct ctcgg 85

<210> 43
 <211> 193
 <212> DNA
 <213> Globodera rostochiensis

<400> 43
 agttcggttc aatgtgctca aggtgatcaa agcatcgggc tcgaagaaag cgttcgacaa 60
 attctgagtc ggccaagcca accgcgaacg gtcatttgtt atgggttccta attgttgctg 120
 tttttcaatt atttgtgtta aatgactgaa tttatgatca acggtatact agtattcttc 180
 tgaaaaagct cga 193

<210> 44
 <211> 219
 <212> DNA
 <213> Globodera rostochiensis

<400> 44
 gaattcattt agatttgttt tgaagctaga aatctttatt ttgggagtca acgacaatgg 60
 gaagacgtcc ggcgcgttgt tctcgtata ttaagaacaa gccgtatccg aagtcgcgct 120
 tttgtcgcgg tgtacccgac ccaaaaattc gcatttttga ttggggtaga aagcgcgcga 180
 ccgttgacga attcccatgc tgcgtgcata tgatatcga 219

AKK110P1

<210> 45
 <211> 489
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 45
 tccgaggcgc ttgaggctgc gcgaatttgt gcgaacaaat atatggtgaa gaattgcgga 60
 aaggacgggt ttcatatgcg cgtcagaatc catccatacc atgtaattcg catcaacaaa 120
 atgttgctct gcgctgggtgc ggaccgtctg cagactggga tgcgtgggtgc gttcggaaaag 180
 cctcagggac tctgtggcgcg tgtcagcatc ggtgatatgc tgaatgacgt gcgtattcgt 240
 gaccaacacc aagctcacgc attggaggcg ttccgtcggg cttaaattcaa gttccctggt 300
 cgtcaataca tctgtcttgc cgcgaagtgg ggcttcacca aattcgatcg cgaggatatac 360
 gagaaatacc gcaaggaggg ccgtgttatc cctgacggtg tgcattgcaa gttactcaag 420
 caacacggac ccgctgaagg agtggctcaa gaacccatt taatcttctg tttgtcttgt 480
 gactcttgg 489

<210> 46
 <211> 101
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 46
 gaattccccg gctcgagccg ggttgacgat gtcctcctcc acctcctctc actgcggttc 60
 gtcctccttc agccggaat tgttcctgtg gctgttgccg g 101

<210> 47
 <211> 485
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 47
 tccaccaaag tccattcgtc gtcgccagtc catttatttc acaaaaagat gattccgctc 60
 tctgtccgat gacgtcgttt ggccaaccgt tgccccctgc accgctttca ctggtgccaa 120
 accgcgcgct ttattttgtg ttcccagaaa acttgccgtt ggagcggccc ttcgacgagc 180
 aaaacgacgg ctccgaggag gaattagccg aagaagcgat gggaacgaag gcgaagaggg 240
 gcgaacggtt cgtccgattc ggcaaaaggc cgcaaacatt tgtgcggttc ggaaagcgtg 300
 cacaacatt tgtacgcctc ggaagggaca cgcaaaggca attcgatggg aaaatgcaaa 360
 gtgaacagca acagaaaaag gcttaaagca aacggcggcg acttttcttt taatgaatgc 420
 gcgccaccg catgacaatt cttttgtgta atgtgttgcg atttttatga tcggtaaatg 480
 taaca 485

<210> 48
 <211> 651
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 48
 atctgttcaa gggactgttc ggcaagaagg aaatgcgcat tctgatgggtt gggttggacg 60
 ctgctggaaa gacgaccatt ctgtacaagt taaagctcgg cgaaattgtc accaccatcc 120
 caacaattgg cttcaacgtg gaaaccgtcg aatacagaaa catctcgttc actgtttggg 180
 acgtgggtgg tcaagacaaa attcgtccac tttggaggca ctacttccag aacacgcaag 240
 gactgatctt cgtcgtggac agcaacgac gcgagcgtgt gggcgaggcg cgtgaagagt 300
 tgaatgcgaat gctggcggag gacgagttgc gcgacgcggt gttgctgggtt ttcgctaaca 360
 aacaggattt gccgaatgcg atgaacgccg ccgaactgac agacagactt ggactgcaca 420
 acttgcgaaa ccgcaattgg tacatccagg ccacctgcgc gacttcgggc gacggactct 480
 acgagggact ggactggctg agcaaccagc tcaagaacag aggctaagct gggttgggtg 540
 ctgttgcaat tgcggcggga attgatgacg attgaattta tttgtgtgtt tgcgcgcgca 600
 gctcttttgt gggacgtccg attaattttg ataattattt tattccgtgt t 651

<210> 49
 <211> 660
 <212> DNA
 <213> *Globodera rostochiensis*

AKK110P1

<400> 49
 gaattcccaa gtttgagatc aattcagttt cacttagaca aaaatgccgc cgaaattcga 60
 cccaactgag atcaaaatcg tgtacctgcg ttgctgcggt ggtgaaattg gtgcaacatc 120
 tgcacttgca ccaaaagtgg gccacttgg attgtcgccc aaaaaaattg gtgaagacat 180
 tgcgaaggcc acacaggact ggaaagggct taagggtacc tgcaagctga caattcagaa 240
 tcgtgtcgcc aagatcgacg ttgtcccatc ggccgcctct ctgatcatca aagagttgcg 300
 cgaacctccg cgagaccgca aaaaagtcaa aaacgtgaag cacaatggca acctgaccat 360
 cgagcaagtg atcaacattg cgcgtcagat gcgccctcgt tcaatcgcac ggaagttgca 420
 gggcaccgtg aaggaaattt tgggaaccgc ccagtcggtt ggctgcacca tcgatggaca 480
 acatccgcac gacattgtgg acgcgatcag agggggagac atcgaaatac ccgaggaata 540
 aagaaaggac ggcgcctccg atttttgtgg gacggacatt gggaatttga ggtgaatgag 600
 ttgccaatth cattcattca tcaattgttg ttattgntgg tacggataaa tttgtaattg 660

<210> 50
 <211> 625
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 50
 gtgccggaac agacgctcga ggaggttagc cgtctgcagc ggacgagctc cttgttggac 60
 gtggcaatcc gggacggcgt cccctacccc ccactgcctc ctacaaaccg atccccgaa 120
 tacatgaaca tgctgaccgg ctcttctctc gtgccaaatt tccgcatcta ctccggcgcc 180
 atcggaccgt acagaccttc gttgcccgtg tacacttaca acacttacca cgggtacttc 240
 ccttaccgca actaccgagg ctacaccttg gcgaatgctt actggtacga ccgatactat 300
 tacttctcgc cgctgtacaa acgaagcatg ttccccaccg gcttcaaaca ttgtgactat 360
 aaagcgaacc cgcactattg gcactacccg cacacctttt gggactatcc ctaccagggc 420
 aaatggttcg actacgacaa ccctcccaat taccggccct actacaacca tcgccttaac 480
 ggatatgctc ggccgtatca ctaccggtcc catgcgctgg cccaccggtt caattaccgc 540
 gaaggaaattg tcaggaaacg ggtctgacaa atcgaactgc tccaaattga cgtggtccgc 600
 attcgaaaga agacgaaaaa agctt 625

<210> 51
 <211> 402
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 51
 gaattccaag tttgagcaac attttgaaaa tgaccgaagc caaaaaactt cccgaggtgc 60
 cggaaacttt gctcaagcga cgcaaaatca gagctgcgca aaaggccgca aaagcaaaga 120
 acaaatgtgag ttctatcaaa aaagcacgga ccaagaagggt ggaaatcttc aaaagagccg 180
 agcagtattt ggtggagtac cgtcagaagc aacgccaat gcttgcgctg aaacgtgaat 240
 cgaagaaagt cggcaattat tatgtgccag aagagcccaa actcgccttt gtggtccgaa 300
 tcaaaggcat caataagatt catccgcgtc ctgcgaagggt tctgcagctt ctccgcttgc 360
 gtcagatcaa caacggcgtt ttcgtaaagt tgaacaaggc ga 402

<210> 52
 <211> 433
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 52
 ccgaccgta catcgcttgg gggtatccga gtcagaagat catccgtcag ttggtctaca 60
 aacgcggtta cgccaaagag aaggacagc gcattccaat aacggataac aacattgttg 120
 agcgcagttt gggcaagcat gacgtgattt gtgtggagga tatgatccat cagatttggg 180
 ccggtcggag cgcacttcaa acaggtgacc aacttcctat ggcctttcaa gctgagcaac 240
 ccggtggggc ggttcaagaa gaagtccaat cacttttgtg gagggaggcg attatggaaa 300
 ccgcgaggac caaatcaaca aattattgga aagaatggtc taatggaagg gaagcggana 360
 aagaaaggaa attgnggcgt ttttctgttg ttgttttgac gataaattgt taactccaaa 420
 aaaaaaaaaa aaa 433

<210> 53
 <211> 768
 <212> DNA

AKK110P1

<213> Globodera rostochiensis

<400> 53

```

gaattcgttt gaggtcaaac tttattagcg tatTTaaca tgtccgaagg aggagcgaaa 60
aagagtagca gCGgtgcca ggggggggtt gatgtcaaga aatttgcat cgatcttgcg 120
tccggtggta ctgCGgCGgc tGtctccaaa actgTtgTtg ctccattga acgtgtcaaa 180
ctcttgTtgC aggtgcaaga tGcttccgct cacatcactg cGacaaaacg ctacaaaggc 240
attattgacg tGcttgTccg tGtgccgaaa gagcagggct ttctgtcact gtggcgTggg 300
aacttgcca acgttatccg ttatttcccg actcaagcgc tgaacttcgc ctTcaaagac 360
acTacaAAC gcatctttac ggagggactg gacaaaaaca agcagttctg gTcgtTcttc 420
gtcatgaatt tggcctctgg aggtgCGgC ggcgCCacgt cGctgacctt tGtttatccg 480
ctgggacttt gcccgtacgc gTttggCCcg tGatgtccg aaaagctggT tcccgcgagT 540
tcaacggttt ggccactgc atcgcaaaaa tcttcaagtc ggacggtccc atcggtcttt 600
accgCGgctt ctTcgTctcc gTccagggca tcatcattta cGcgCGcGcc tactttggat 660
gctttgacac cGcgaagatg attttcGcgC cggatggcaa gcagatgaat ttcttctca 720
catgggcat cGctcaggtc gTcaccgtgt cgtccggtgt cctctcct 768

```

<210> 54

<211> 338

<212> DNA

<213> Globodera rostochiensis

<400> 54

```

gaattccagc agattaattg gaatggctga gaacatcgaa gagattcttg ccgaaatcga 60
cggctcccaa attgaggagt atcaacgctt tttcgacatg ttcgaccgcg gaaagaatgg 120
ttacattatg gccacccaaa ttggacaaat tatgaacgcg atggagcagg actttgacga 180
aaagaccctc cGaaaattga tccgcaagtt cGacgCGgac ggttccggca aactggagtt 240
cGacgagttc tGcgcgTtg tGtacacggt ggccaacact gTggacaagg acactctgcg 300
aaaggagctg aaggaggcat tccgactctt tgacaagg 338

```

<210> 55

<211> 267

<212> DNA

<213> Globodera rostochiensis

<400> 55

```

gaaattgcgc ccgatctcag cgacaaggat ttggaggcgg cggtcgacga aattgacgag 60
gacggcagcg ggaagatcga attcgaggag ttctgggagt tgatggcggg cGaaaccgac 120
tgagaaaaga gcaaatcgat ccaaatccaa acggaccctg cccatttcac ctccatccgt 180
ccgtcgTatt attatatTTt ccagtggaaT tttccatta aaattcggTg aaagtaaaat 240
aatttgacga aaaaaaaaaa aaaaaaa 267

```

<210> 56

<211> 597

<212> DNA

<213> Globodera rostochiensis

<400> 56

```

gaattcgctg gacacttcgc atccggagta cagccacgag cagagcatcg accagaccag 60
catcccctac cagatgggtt cgaacaagta cgcctcgag aagggcatga ccggctttg 120
acagcccctg tgggaggTgc ttgaccgctc catctcgtac cagaaccgca agtcgcaagg 180
aatggTtcgt ctacagtcggt gtaccaaccg gttcgctcc caggcgggca tgaccggctt 240
cggcacaccc aggaacacca cctatgaggc ggaggcaggc gagctgccct acgaggacat 300
gaagaagtcg gaggcgatca tcccgTccca ggccggttg aacaagggcg actcgCagaa 360
gttgatgacc aacttcggca cgccccgtaa caccaccacc aaggtcaaaG tggagaattt 420
ggcggaaaatt ccggaggaca ttttgctgaa aggacacggc gaggtgcgcc tgcagtccg 480
taccacccgg ttcgctgcc agaagggtt cgtcgcgTtC ggtaccggac gtgacgtgtg 540
ccgtga9999 gtgaacgtga acgtgctgcc gggcgacttg gagccgcttc cgaggga 597

```

<210> 57

<211> 80

<212> DNA

<213> Globodera rostochiensis

AKK110P1

<400> 57
 ggcattgtgc gtctgcaagc cggtagcaac aagttcgact cgcagaaggg catgaccctt 60
 ttcggtacgg gcccgtcgtg 80

<210> 58
 <211> 513
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 58
 gaattcgcca caccgctcac atcgcggtgca aattcgccga acttaaagag aagggtggacc 60
 gncggtctgg caagaaagt gaggacaacc cgaagtcgct gaagactggc gacgccggaa 120
 ttgtcgaact gattccgacc aagccgatgt gtgtggaggc attcactgac tacgcaccgc 180
 tcggccggtt tgctgttcgc gacatgaggc anactgttgc cgtgggcgcg atcaaatacag 240
 tggagaagac ggaaggcggg ggcaaagtga ccaagccagc gcagaaggtc ggcgcgactg 300
 gtggcgggaa gaagacatga ccaaggggag gggcggttcc ctaaggcca accgtcgacg 360
 aaaaatgcgac caacctcttg tttatcgttg tcttattcag ttccttccac ccgtctctat 420
 ccatattgtc gttgcgttgg ataattgttt atttttgtt attgtcctgg ttggaaaata 480
 aatttggtca attaaaaaaa aactcgtgcc gaa 513

<210> 59
 <211> 393
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 59
 gaattcggtt gagcgaaaaa aacatactat acaatggcaa caactgagaa gcctcaggtg 60
 gttcaacagc ccgtgcaggc ctttggccga aagaagacag caacagccgt tgcgtttgca 120
 aaaaggggca agggcttgat caagggtcaat gggcgctcct tggactacat gcagccggag 180
 attctgcgca ttaagctcca ggagccaatt ctcatgttgg ggaaggacaa atttgagggg 240
 atcgacatac gaattccgct caagggcggg ggacacattg cgcaaattta tgcaattcgc 300
 caagcactgg ccaaggcact ggtcgctttc taccagaaga atgtcgacga gcagagcaaa 360
 aaggaactga aggagcaatt tgttgcttac gac 393

<210> 60
 <211> 154
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 60
 cagagccaa agaaattcgg tggacccggg agctcgcgct cgctaccaga atcgtagcgt 60
 taagaaataa tttttagat caaatgtttt gatgatgatc cttgtttttg ttgttgataa 120
 aaaaaattta taaaaaaaaa ccgccgatac tgac 154

<210> 61
 <211> 666
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 61
 gtattccaag tttgagcgat cagagttcct caatctatta tcaactgttt tccatcaacc 60
 aactgtcatc atgcaaattt tcgtcaagac gtcaccggc aagaccatca ctctcgaggt 120
 cgaggctagc gataccatcg agaacgtgaa agccaagatc caggacaagg agggcattcc 180
 gcctgatcag cagcgtctga tcttcgccgg aaaacagctt gaagacggac gcaccttggc 240
 cgactacaac atccagaagg agtccactct ccactctcgt ctgctctcc gtggcggaat 300
 gcaaattttc gtcaagacgc tcaccggcaa gaccatcact ttggagggtc agggcagcga 360
 caccatcgag aacgtgaagg ccaagatcca ggacaaggag ggcattccgc ctgatcagca 420
 gcgtctgatc ttcgccggaa aacagctcga agacgggcgc actctggccg actacaacat 480
 ccagaaggag tccactctcc atctcgtctt gcgtcttcgt ggaggagaga actgaatcgc 540
 gggctgatgg aaagatgacg aatatgatgt ctattcgatg acttgtctct ttcgatataa 600
 ttgatttgtt tccatttgtc ggtcatcaaa tctttatgac cccctcattg ggcattggaac 660
 gataaa 666

AKK110P1

<210> 62
 <211> 213
 <212> DNA
 <213> Globodera rostochiensis

<400> 62
 gaattcgttt gagaaacttt ttcaaccatt cattcaaag tctcatcaag tgacacgggc 60
 agcactcaac cacgggacgc gtgtactgag cgtgttggag aagggtcaagt tgggtctgctg 120
 gtttgaggag acacattcgt tcgcgcaagt ggctcgaaga taccgggcag aatttggtat 180
 ggaaccaccg cagttggacc aagtgaagaa gtt 213

<210> 63
 <211> 488
 <212> DNA
 <213> Globodera rostochiensis

<400> 63
 agcaccggct caatcctcaa tggcacaacg acggcattct cggcatagag agacggagtc 60
 ggtcttggag aacaacagcc aattcccgtc gtaagcgatg cgggactgga tgcggaagaa 120
 cagctgagaa tggccagaat gtgagccgga ggacctgaag atttatgaac gaaattttcc 180
 agtgaagtgg accaacgctc ttcgacttta tctgctttgt gtaaagtgtg tagaatcggc 240
 ttccaattca aaggcttttc attccccaac ttttattttt gcgcaaaaaa tttcttagga 300
 taagcgtgaa taattttattg atttgttttt tctttctttt atctccgcct cgaagtgcga 360
 agtggttcctt ttggcccgtt cccttttgtt ttgaatgtta ttccattccc atcccctcac 420
 tttctcatat ttgtgacatt cagctgcatt gttcgactcc catttaaaag ttgagtgaag 480
 tgcgattg 488

<210> 64
 <211> 249
 <212> DNA
 <213> Globodera rostochiensis

<400> 64
 wccrgakbng aacahcdkdg vhwatnvcbn gschvbwagc rngtcsvddb wgnhnsswtg 60
 gkgdyrbwnt msnwrmanrg artsstsgaa ttcccaagtt tgagagtaaa tattattagc 120
 taaaaatggc agtcggaaag aataagagaa tgggcaaaaa gggagccaag aagaaggctg 180
 tcgatccgtt cacacgcaa gaatggtacg acatcaaagc gccggcgatg ttcaacacatc 240
 gaaatssts 249

<210> 65
 <211> 362
 <212> DNA
 <213> Globodera rostochiensis

<400> 65
 wcbcrbhdby ytsgcrsnck tbdshbcysy gcdwkmtnvk hscngdckty nykklkvbmr 60
 ntmsnwrman rgartsstsg tcaaccgtac tcagggaacg cgcatttcga gcgactttct 120
 aaaaggccgc gtttacgaag tgtcactggg tgaccttaac agcactgacg ccgactttcg 180
 aaagttccgc ctgatctgtg aagaggtaca gggcaagatt tgcctgacca actttcacgg 240
 aatgtcgttc actcgggaca aactgtgctc tattgtcaag aagtggcaca cgctcattga 300
 ggcgaatgtg gcagtgaaga ctaccgacgg tttcatgctc cgactctttt gtatcgggtss 360
 ts 362

<210> 66
 <211> 128
 <212> DNA
 <213> Globodera rostochiensis

<400> 66
 aatcaaatta agaagacgag ctatgcaaaa gcctctcagg tgcggatgat tcgtgccaag 60
 atggtggaga tcatgcagaa agaggctctt tccggcgatc ttgaangaaa gtagtcaaca 120
 agcctgat 128

AKK110P1

<210> 67
 <211> 502
 <212> DNA
 <213> *Globodera rostochiensis*

<400> 67
 gaattccatt aaaaaactaa acgaacaaat ctaaagatgg ccaccgaagt ggaggaaaat 60
 gttcctacgg ttgacccatg ggggtgctgt gaggaagtgg gtggtgaaga gtcgatgcag 120
 ttggtcagcc ttgacgttac cgagggtcaaa ctgttcggaa aatgggtccct taacgatgtg 180
 gaagtgtccg acatttcgct tgtggattat attgcggtga aggaaaaggc ggccaaatat 240
 ctgccgcaca gcgccggccg ttaccaacag aagcgccttc gcaaggccac ctgtccggtg 300
 gtggaacggt tgtctttgtc aatgatgatg cacgggcgga acaacggaaa gaaactaatg 360
 gcggtgcgca ttgtgaaaca ccccttcgag atcatcacct gctaccggag agaaccag 420
 ccaagtgttg gtcaatgctg tgataaacag tgggccccnc gaagattnca cacgtatcgg 480
 acgtgcgggc actgttcgtc ga 502

<210> 68
 <211> 519
 <212> DNA
 <213> *Meloidogyne incognita*

<400> 68
 gcaaaacttt atcaaataaa aaatttatat ttgccaaaca aatttatgaa taaaaattca 60
 ttaatcatta aaactacatt taaaatatac tttttagaga atgtcgtcta aaatattctt 120
 ttctcccctt tatgcatcta tctaaccaga cttggaagca atatggctaa tcaagtcaac 180
 aatacggcag gaatacccaa actcgttatc ataccagcta accaatttaa caaatgcgg 240
 gttgagaacc ataagagcct cggcgtcgaa aatagacgaa tgagtgtcgc caagaaagtc 300
 ggtagaaaca acctggtcct cagtatatcc aagaatccct ttaagctttc cttccgaagc 360
 agtcttaatt gcattcttaa tagcctcctt cgttgctggc ttctccaaac gagcagtc 420
 atcaacaacg aaaacgtttg ggcgtcggca cacgaaaagc catttcgggt aagcttccca 480
 tccaattcat ggattgacct ttccaacagc ctttgagc 519

<210> 69
 <211> 218
 <212> DNA
 <213> *Meloidogyne incognita*

<400> 69
 ttgattcttt attagtggac aatgacggaa gaccagaaga agttgccgat ggtgcctgag 60
 actgttttga agcgaaggaa agttagggct gctcagcgtg cttctctact caagaataaa 120
 ttggagaata ttaagaaggc taaggttaaa acgcaagtta tctttaaacg tgctgagcaa 180
 tacttgattg catatcgacg taagcaaaaag caagagtt 218

<210> 70
 <211> 293
 <212> DNA
 <213> *Meloidogyne incognita*

<400> 70
 taagaaagca ggggaatttt atgtcccaga tgaacctaaa cttgcttttg ttgtgcgtat 60
 taagggaatc aacaaggtta atttaaattt gctataaagt ttaggatggg tttagacaat 120
 tcttctcttt taatgctttc taactttttc aaaaaagtta tgattttatc acccattaat 180
 ctacaaattc tttaatttat cagatccatc ctgcgtcctg aaaagttctt caacttttcc 240
 gcttgctgca aatcaacaat ggagttttca ttaaattgaa taaagctaca atc 293

<210> 71
 <211> 422
 <212> DNA
 <213> *Meloidogyne incognita*

<400> 71
 aatgcaatta agactgcttc ggaaggaaaag cttaaaggga ttcttgata tactgaggac 60
 caggttgttt ctaccgactt tcttggcgac actcattcgt ctattttcga cgccgaggcg 120
 taagttttga ttttctaaga ttatatttaa ctttttaatt ttttcagtct tatgggtctc 180

AKK110P1

```

aaccgcatt ttgttaaatt ggtagctgg tatgataacg agtttgggta ttcctgccgt 240
attgttgact tgattagcca tattgcttcc aagtctgggt agatagatgc ataaagggga 300
gaaaagaata ttttagacga cattctctaa aaagtatatt ttaaattgtag ttttaatgat 360
taatgaattt ttattcataa atttgtttgg caaatataaa ttttttattt gataaaagtt 420
tg 422

```

<210> 72
 <211> 374
 <212> DNA
 <213> *Meloidogyne incognita*

```

<400> 72
atctgagcat aaggaaactt ggcctcaagc tatagagcag accgattatg tggcaccgac 60
tgagccagtt aaactggact tcaacgttcc gcttattagt gattgggctg ctgcttctga 120
gtggcctcaa gaagaggaag ctcagggttc acctactgca ccaattgggtc agccacagcc 180
tcaacagcag caaactcaac aaggaggtga ttggaactct ggtactagt gatggtgaag 240
ggcaggaaaa ttgatagaaa gagaaattat tatggaataa atgtaataca tgttgtgtgc 300
tgattttatt gttacatata caacaagttt tatittgttg tttatttaat aaaagtgtgt 360
aattaaaaaa aaaa 374

```

<210> 73
 <211> 120
 <212> DNA
 <213> *Meloidogyne incognita*

```

<400> 73
tttttttttt tttttcttca tcaatatttt gaagtgaaga accagaagta gttgcattcg 60
agctttcaaa ttttgttttt tgattactct ttaaacaaga ttcaactgat ggatctactg 120

```

<210> 74
 <211> 369
 <212> DNA
 <213> *Meloidogyne incognita*

```

<400> 74
gtctaaccac tctagagcta ttcggttcgt ctgtctgttg attattagat gttgattgaa 60
cagcactagt ctctgatgta gttttcttca atctcatttt taagtgatgt agaggaagtt 120
tagaattctg attgctatcg tcttctttct ctctctttta tggctttttc aatttatctt 180
cttctttttc ttgtccattc ttttcttcat tcttttcaaa aggttcagga aattttaatt 240
cagaccgctg ccttttaact gctgtatcta aagaaaaccc tctaggcaac gtcccagttc 300
cactcaaatt caattttgtt aaatttttgc cagatctaag tccttcttcc ttttgaacga 360
attgaactg 369

```

<210> 75
 <211> 529
 <212> DNA
 <213> *Meloidogyne incognita*

```

<400> 75
ttttgttttt tttttttttt ttatcagaaa aaagttttaat cagaaaaaaa aattaaaaca 60
aatctaaata aggctctatt ctaagtttat atttttcttt tacataaacc gtcaaccctc 120
caagtttttc aatgcttggg ggttttaatg gatcctctgg taataatttg taggctagaa 180
aaaagtgtgc agcaaaaagg aaaagcatca ttcttgctaa ggcttctcca gcacattgcc 240
ttttcccccac accaaaagct attagctcgt cagctttttt taatttccct tcattgtcta 300
tataacgttc agggtcacaaa ttttggggat ttgggtatat ctttggatca aaaagaacat 360
ccgatacttg gggatcaca aatgtacct taggcaacac aaactttcca acattcaaat 420
cttccaaggc taaatgcccc aaattgaaag ggactaaatt aacgagtctt aatgtttcat 480
taacaacagc atttgtataa attaatatag gtctgtgttc caaactaat 529

```

<210> 76
 <211> 449
 <212> DNA
 <213> *Meloidogyne incognita*

AKK110P1

<400> 76
 agtttttttt tttgaataaa agactttttt ttattataaat ggcttcgcaa actgcaggaa 60
 ttcaacaatt acttgcagca gaaaagcgtg ctgcagaaaa gattaatgag gcacgtaaaa 120
 gaaaggcaca acgacttaaa caagcaaaac aggaagcgca agctgaaatt gacaaatata 180
 gagaggaacg tgaaaaaacgt tttaaagagt ttgaacataa ttacctcggc gctagagatg 240
 atatttctgc acaataaaag cgtgaaactg atgagacgct taatgaaatg actcgtagtg 300
 ttgctgctaa taaacagcag gtaattgttc gtctacttca acttgtctgt gacattcgtc 360
 cagaactgca tcacaattta caacttcaac ttaagcttaa tgaaaagcct gcctaatttg 420
 tagttgattg attataaaaa tgaaattga 449

<210> 77
 <211> 643
 <212> DNA
 <213> Meloidogyne incognita

<400> 77
 atttataattt gaacaaataa ttttaacaaa aagtattggct cgaggaccac agaagcattt 60
 gaagcgtttt gccgctccaa agaattggat gttggacaaa ttgggtggag tttttgcccc 120
 acgtcccatg tgcgggcctc acaagcttcg tgaatcgctt cctcttattt tgtttcttcg 180
 taatcgtcta aaatatgcac aatcttataa tgaagctagg atgatttgca aacaacgtct 240
 cattaaagtt gatggcaagg tgcgtacaga aatgcgcttt ccagctggat ttatggatgt 300
 ggtttccatt gagaaaactg gcgaagtctt tcgtcttctc tatgatgtca aaggacgttt 360
 cattactcat cgcatacaaa aggaagaagg tcagcttaaa ttgtgcaagg tagtaaagca 420
 agcgattggg ccaaaacaag ttccttatat tgttactcat gatgcccgta ctattcgtca 480
 tccggatcca cacatcaagg ttgacgacac tgttgctgtt gatataaaca ctggaaaggt 540
 tacagatcac attagatttg attctggtta tgtttgtatg attactggtg gtcacaacat 600
 gggacgtgtt ggtattggtt gacatcgtga acgccaccct ggt 643

<210> 78
 <211> 584
 <212> DNA
 <213> Meloidogyne incognita

<400> 78
 atttcctcta aaaatgaatt taaaagaaca acaaatatat ttaaatattc aattattatt 60
 ttttattttt gctgtcagta gttttttgac aactaaggga agtgaagtaa aacaacgaga 120
 aaataataaa ttggaatata ataaaaatga aattgagagg caaaaagagc aattaattcg 180
 agatttgatt gccctcttaa cacgtgaaag gcaatattca cgagattggc aacaatcaca 240
 acagcaacaa aatttcatta acagtttttg cccttcccca catttattcc cctcttcagg 300
 cattgaatgg ccccaacaac aacaaaaaat atttttggaa gaaggggaag tagaagaacc 360
 tttagaggaa aatgagaagg aaaaaagagc acaaaccttt gttcgtttcg gaaagagagc 420
 acaaacattt gttcgttttg gaaaaagggg acagactttt gttcgatttg ggagagattc 480
 aaaacatcaa cataacttgt cagatcagaa gcagttaaaa actgacaaac aataaaaatg 540
 atgaattatt taaaaatttt tttaatgatc ttttaattaa aatt 584

<210> 79
 <211> 556
 <212> DNA
 <213> Meloidogyne incognita

<400> 79
 atcaagcatt aaatatgcag atttttgtaa agactctcac cggaaaaact attactctcg 60
 aggttgaggc ttctgatacc attgagaatg ttaaggcaaa aattcaagat aaagagggtg 120
 tcccgcctga tcaacagcgt ttgatctttg ctggttaagca acttgaagat ggacgaacct 180
 tggctgatta taacatccaa aaggagtcta cacttcactt agttttacgt cttcgtgggtg 240
 gaaagggttc cggttcattg gctcgtgctg gaaaagggtc tgctcaaaact cctaagggtc 300
 aaaagcagga acataagaaa aagaagcgcg gccgtgcttt ccgtcgcatt caatataacc 360
 gtcgcttcac caatgttgct acttctgggg cgggacgccg tcgtggccct aactccaacg 420
 ctgcataaga gaatggctgt atcttgatga atgtatggtg atataatcaa ttttaatacat 480
 tcgactntat gaagttttct gttattcaag ataaatcttt ttgttgaaaa aaaaaccaag 540
 tttgagatca gttact 556

<210> 80

AKK110P1

<211> 424
 <212> DNA
 <213> Meloidogyne incognita

<400> 80
 aacattggtt taattaaaat ttaccctcc ttagcaatg acatcagaca gacttggccc 60
 agtagttcca gatttgacag cccaagagac caacagactt gaacgaacta gttctttggt 120
 cgatttggca attcgggatg gagttccata tccacctagg cctgcaatta ataatgttcc 180
 tccatacctg aatatgttga ctggaacggt ttctgtacca aatgtaaatc agtacacggg 240
 tgcaataggc cttatcgac cagcaaatcc tgtttatact tattatagct ataaatgcta 300
 ttttccgtat agaaattatc gaggtacac actgacggat gcttactggt acgaccgtta 360
 ttattatatt tcgccaatat acaaacggtc aatgttccca attagattcc ggcattctga 420
 ctac 424

<210> 81
 <211> 89
 <212> DNA
 <213> Meloidogyne incognita

<400> 81
 attatccaca cacctattgg agctaccctt accaaggaaa atggtacgac tatgacaatc 60
 caacanatta cgcgccattc ttgaccca 89

<210> 82
 <211> 168
 <212> DNA
 <213> Meloidogyne incognita

<400> 82
 tttttttttt taaaatttat tcattaacaa atgaccttaa cagataaaaac ttaacagtca 60
 aaagacaaca taattttcaa ctttttcaat attatccttt ttaacgggtt gattttgcaa 120
 ctcgctccaa ttcgtccttc ttcttgatag catatgaatt gctcgaac 168

<210> 83
 <211> 67
 <212> DNA
 <213> Meloidogyne incognita

<400> 83
 aattcatcag ccagacattc agcaattggt ttgatattac ggaaagaagc ttcacgagac 60
 ccagtac 67

<210> 84
 <211> 42
 <212> DNA
 <213> Meloidogyne incognita

<400> 84
 taacacgacg aagaggcgaa acatcaacag cctgacgacg aa 42

<210> 85
 <211> 429
 <212> DNA
 <213> Meloidogyne incognita

<400> 85
 tatacgagta gaatcctccc gtggtcctcc attaataaca gcgccaacaa gtatttgaac 60
 tggattctct ccagtcaaaa tatgtataat ttcaaaagcg tgcttcacaa tccgaacagc 120
 catcaacttt ttaccattgt tacgtccatg catcatcatc gaacaaacca aacgttcaac 180
 aatcggacaa tgagcctttc gaaaacgttt gatttgatat cgaccagcac tgtgcggaac 240
 atatttggcc gatitgtctt taacagcaat ataatccact aaagaagcat cattaacttc 300
 gatatcgctt aaagaccatt taccaaacaa tttaatttca ggaaaatcaa ttgtagtcac 360
 ttgcatatcc ccttgtccac caggaacatc agttgcgccc caattatcat cagcgggtaa 420

accatctcc

AKK110P1

429

<210> 86

<211> 435

<212> DNA

<213> Meloidogyne incognita

<400> 86

tttgagtttt	taaaaagtac	atactattta	atttttaaca	aattattttg	atcaatttta	60
aatttttctt	tcatcatttt	ttaattttaa	aaacatttta	acaaattaca	agaacaacaa	120
acataattgt	ctctttttta	ttataaaaatt	taaagtttaa	taagttttta	aacatttctg	180
actggagtag	gtgtacttag	tgtttttagaa	aaggcaaaat	tagtttggtg	gtttgaagag	240
acaaattctt	ttgcacaagt	agcgagaaga	tatcgagcag	aatttggaat	ggaaccccca	300
catatggatt	tagttaaaaa	attacatcaa	cgttttctca	atactgggtc	tgttttctaat	360
ggaaatactg	aacattttga	agttaatcca	acaatggaaa	catcgacatc	ctcaacagag	420
ggtgtagcag	atccg					435

<210> 87

<211> 501

<212> DNA

<213> Meloidogyne incognita

<400> 87

gttttttttt	ttttttttta	aacaaaatat	cgagtcttta	taagacaaaa	ataaaagaca	60
aaagcaattt	agttttatca	ataaaaattaa	aaatagtcaa	tgtctcgttt	cactcattag	120
atttgtggcc	ctaaagaggg	ccgtttgggt	ttggttggtg	tacttcagct	gccttccacc	180
aattgttcct	tagccacca	atccgtaaag	agtacgtcct	tggcgtttca	acgcatagac	240
gacgtccatg	gctgtgaccg	tctttctctt	ggcgtgtacg	caataagtta	ccgcgtcgcg	300
gatcacattt	tcaaggaaga	ctttcagaac	acctcgagtc	tcctcgtaaa	tgagcccggg	360
aatacgtttc	actccaccac	gacgtgccaa	tcgccggatt	gccggtttgg	tgataccttg	420
gatgatata	cgcaagactt	ttcgggtggc	cttagcgcct	ccctttccaa	gtcccttttc	480
gccttttact	cgtccggaca	t				501

<210> 88

<211> 270

<212> DNA

<213> Meloidogyne incognita

<400> 88

ggaagtgtgt	ttaagataaa	tgatgatta	gaaataaaaa	tgaattgatt	aaaaattacg	60
ttagaataat	aatggaatat	ataaaaaata	attggatgat	ttaataaaaa	aaaaaaagag	120
agaactagtc	tcgagttttt	tttttttttt	tttttaanaa	ttaacaattt	atctcatttt	180
cctcttccat	gaaaattaac	aaaaagacga	caacttaatc	ccataattaa	catcattttt	240
aagcttcagt	cggcatgctt	cgaataatgt				270

<210> 89

<211> 286

<212> DNA

<213> Meloidogyne incognita

<400> 89

caagcgggtc	ccaactcaat	gttgttgcca	tgatactcgt	gaacaccagt	tctcgccaac	60
atagaatagt	actcaatctc	actgcgtcta	aggcttggag	tattattcga	aataataaca	120
agtttagcct	ttccagaacg	aagagtcctc	aacgtctgct	tgtagcccaa	acaatacttg	180
cccgatttgg	taaccatggc	gagacgagca	ttgatatttt	ctgtggactt	tttctgtttt	240
ccaacaacca	ttgtaacgca	aaattaaaaa	ctctttttta	acaaat		286

<210> 90

<211> 391

<212> DNA

<213> Meloidogyne incognita

<400> 90

AKK110P1

agatatgaca	tcagacagac	ttggcccagt	agttccagat	ttgaccagcc	aagagaccaa	60
tagacttgaa	cgaactagtt	ctttgggtga	tttagcaatt	cgggatggag	ttccatatcc	120
tcctaggcct	gcaattaaca	atgttcctcc	atacctgaat	atgttgactc	gaacattttc	180
tgtaccaa	gtaaatcagt	acacgggtgc	aataggctct	tatcgaccag	taaatcctgt	240
ctatacttat	tatagctata	aatgctat	tccgtataga	aactatcgag	gctacacatt	300
gacggatgct	tattgggtacg	accgttatta	ttatttttcg	cctatataca	aacgggtcaat	360
gtttccaatt	agattccggc	actctgacta	c			391

<210> 91
 <211> 131
 <212> DNA
 <213> Meloidogyne incognita

<400> 91
 attatccaca cacctattgg agctaccctt accaaggaaa atgggtatgac tatgataatc 60
 caacaaatta ccgcccgttc ttcgaccac gcacacgcgc atcattttca agaccttatg 120
 attacacatc a 131

<210> 92
 <211> 571
 <212> DNA
 <213> Meloidogyne incognita

<400> 92
 ttggtgacgac aacaaaaaaa ttttattttat tttttaacaa cagaaaaata tactttttta 60
 tttttaatat ttttccatga ttcaacagcc atacttttct cattttaata cttcttaaac 120
 cctcaaaaaa ttcattttatt gacgaccagc agcagggtgt tgctgctgtt gttgaccacc 180
 acccccttgc gcttgacctt gctgttgctg tcccttcacg tcaacaggca aattgagttg 240
 caaataatca accatctcct tagtctcttg atcaacacta atagttggat gttgagaagc 300
 atcaagatag gaaacttctg gaacccaatt atcacgacgc tcacgctctt cttcttgcaa 360
 tttaatagaa attccacgaa caggctcttt ttcgatacgc ttcacaaat gggtataaaa 420
 accagcaatt tgattacgca tccgtttgct aggaataaca gcaatttcct cacaaattcg 480
 tttgttcaca tgaaaatcat aagtcaagcg tgtataatat ttgtcaataa taacacgaga 540
 tgctttcttg acagttttga gagaaccgat t 571

<210> 93
 <211> 671
 <212> DNA
 <213> Meloidogyne incognita

<400> 93
 ttgagaatt taacttttct aaccaaact tttatttttg tctttgatgt ctactcaagt 60
 accaatacgc gtgctggtta ctggagcagc tgggtcagatt gggtattctt tgggtattca 120
 aattgcaaag ggtgatgttt ttggaaagga aacgcccatt gttctggtta tgggtgatat 180
 tcctccaatg gccgaagtgc tttaaaggagt ggaacttgaa ctttacgatt gtgccttggc 240
 gaattcttata gctgtcgagc cagtcacgac tgaagaggca gcgttcaaag acattgatta 300
 tgcttttctt gttggtgcaa tgcctcgaaa ggaagggaatg gaacgaaagg atttacttgc 360
 tgctaattgtg aaaatatatta aatcgcaagg attggctcta gcaaaatatt caaagccaac 420
 tgtaaggtt ctggttggtg gaaatccagc aaatacaaat gcttttattt gtgcaaaata 480
 cgcagcagat aaaattccag caaagaatgt cagcgcctatg actcgtcttg accataaccg 540
 tgcaattgcc caaatagctg ctcgttggtg gggtgactgt ggatctgtga agaaagttat 600
 aatttgggga aatcattcaa gtacccaatt tcctgatgtt aaacatgcta aagtaattaa 660
 aggtggcacg g 671

<210> 94
 <211> 289
 <212> DNA
 <213> Meloidogyne incognita

<400> 94
 ggctgtaaat gatgtgccgt ggatacagaa tgaattttatt tcgaccgtcc aaaagcgcg 60
 agctgttatt atcgaaaaac gcaaactgtc cagcgcaatg tcggcagcaa aggcggcatg 120
 tgatcacatt catgattggc acttttgaac aaaagatggc gattgggttt ctatggccgt 180
 tccttccgat ggttcttatg gaattccgga aggtttgatc ttctcatttc caattacaat 240

AKK110P1

tgatgcanaa acgcgtgact ggaaaattgt acaaagatta gaactcgat 289

<210> 95
 <211> 262
 <212> DNA
 <213> Meloidogyne incognita

<400> 95
 aattttaactt ttctaaccac aactttttatt tttgtctttg atgtctactc aagtaccgat 60
 acgcgtgctg gttactggag cagctgggtca gattgggtat tctttgggta ttcaaattgc 120
 aaagggagat gttttcggga aagaaacgcc catcgttctg gtaatggttg atattcctcc 180
 aatggccgaa gtgcttaaag gagtggaaact tgaactttac gattgtgcct tggcaaatct 240
 tatagctgtc gagccagtca cg 262

<210> 96
 <211> 323
 <212> DNA
 <213> Meloidogyne incognita

<400> 96
 aagacattga ctatgctttt cttgtttggtg caatgcctcg aaaagaagga atggaacgaa 60
 aggattttact tgctgctaatt gtaaaaatat ttaaatcgca aggactggct ctagcgaaat 120
 attcaaagcc aactgttaag gttctggttg ttggaaatcc agcagatata aatgctttta 180
 tttgtgcaaa atatgcagca gaaaaaattc cgacaaagaa tttcagcgct atgactcgtc 240
 ttgaccataa ccgtgcaatt gcccaaatac ctgctcgttg tgtgggttgac tgtgggtctg 300
 tcaagatagt tataatgttg gga 323

<210> 97
 <211> 717
 <212> DNA
 <213> Meloidogyne incognita

<400> 97
 aatattttta acaaacgatg taacagaaaa acaaagtttt ttttaacaaat tttcttgaac 60
 cttatttttt ttcaaaacat ttttttattt aaattttaaac ctctcttcat ttctcttaaa 120
 cacttttcctg aactggagggt tcataagcat ctggacgact ttcaataact tctccacttg 180
 ctgtagttat agcaacttgt ccaccaccac ttccagcacc ctctccatgc atatccaaaa 240
 gttttccaag ttcaaatttt ggttttttca aaatttttac ttttcgaata taaacgtctt 300
 gaagtggata gaaataagaa caagactttt caatgtcttt tccaatagaa tcaggaatta 360
 atttgctgac aacttcttta agatcgcatg aagaaacctc gcgatgaata atctcaacca 420
 tcctagcacg aatttgacgc acttgagacg attttgcata actagtcttt ttcatctggg 480
 ttggagcttt ctttgtgaag ccaatacgaag acaatcgaag caaataacca tcagttgttt 540
 tgacagcaac atttgcttca attaaagtat gccacttttt gacaatagaa caaagcttgt 600
 ctgagataaa agtcattcca tggaaattgg tcaaacaacac tttgccttga acctcttcac 660
 aaataagtcg aaatttgcca aagtcagctt cgggtgttgtt cagatcacca agagaaa 717

<210> 98
 <211> 758
 <212> DNA
 <213> Meloidogyne incognita

<400> 98
 gacaagttta accttgtgtg actttatcta tattcttgtc taaataattc taacaaattg 60
 taacaacaaa caaaaatggg cgagcaagac aaaaagaaaag ctggcggcgg cgatggtggc 120
 aaaaagaagg atggcttcga tgccaaaaag tttgcgattg atttggcttc tggaggaact 180
 gccgctgcgg ttctaaagac ggctgtggcg cctattgaac gtgtcaagtt gttgctacag 240
 gttcaagacg cttctcagca catcgtctgc gataaacgct ataaaggaaat aattgatgtg 300
 cttgttctgt tgcccaaaga acagggagtc cttgcttttt ggctgtgtaa tttggctaac 360
 gtgatccgtt actttccaac gcaagctctc aactttgcgt tcaaggacac ttacaagagg 420
 atctttcatg aaggtgttga caagaacaaa cagtttggca aattcttttt gatgatgtg 480
 tttatgagca aaaaatttcct tggttggaat agacctaaaca gttgaagagt atcttgcctt 540
 ctgtgatacg tatacaacac tctcttcaat tggagattca atgttgagtg gagatgtctg 600
 tagtaatcct ctgttacaat cacttaacaa ctcaatcaat tccaatgcca ctgctcagaa 660
 ttataactcc tcaacaattg gccgaagcta aaaactacgt ttcaacatgc tacagctact 720

AKK110P1
tcaaaatcga aacagattgt tttaaacgtt tgaaattt 758

<210> 99
<211> 154
<212> DNA
<213> Meloidogyne incognita

<400> 99
ttgagttcgt tggcacattt gttgtgttac aaaacgaaaa ttattgggaa cgggtttcag 60
tgccatttct cgcaggttat tggcacttca cacatttgta ccaataacaa cgttaccgtt 120
tataatcaaa ctgttcctca aagttatgcc catt 154

<210> 100
<211> 125
<212> DNA
<213> Meloidogyne incognita

<400> 100
ttcagaatac tcaaggtctt atattcgttg ttgatagtaa cgacaaagag cgtattgttg 60
aagctcgtga ggaattgatg cgtatgttgt ctgaagacga acttcgcgat tctgtactcc 120
tcgta 125

<210> 101
<211> 219
<212> DNA
<213> Meloidogyne incognita

<400> 101
cttgccgaat gctatgaacg ctgctgaact tacagacaaa cttggacttc acacgctgag 60
aaatcgtaac tggatatatcc aggctacttg tgccacttca ggagatgggt tgtatgaagg 120
tttggactgg ttgagtaacc aattgaagaa tcaaggttaa atgagtctaa ataaaaatgg 180
agagggggaaa gaggagaggt taatttttta aggaaaaaa 219

<210> 102
<211> 473
<212> DNA
<213> Meloidogyne incognita

<400> 102
gttttttttt ttttttttta aattccaagt tttctttcaa atgagagaat agggagaatg 60
atgggggaaa aaataggagc aagccaaaaa gccaaaaaaa aatttttttt ttaaattgatt 120
tttgtaaatg tgtgaaaagg tgtgtgtcaa ttgtagagtc aaatgtcgtt gccttccttc 180
cactaaaatt tctctttcct ttcttttctc ttctaaaatt ccttcaaagt cgatccaacg 240
aaatttcagc ctcctctgga tattccaact cccaaatagc cttcaaagt ttgcctttta 300
cgtcacgagg agtaccaa atccagtcac caactttgaga gtctccctta ttccaaccgg 360
cctgggatgg aattatcgtt tctgacttct tcatatcttc atatggaagt tcgccagact 420
ccgcctcgtat tgttggtgtc cttggcggtc caaacctgt catgcccgct tgc 473

<210> 103
<211> 114
<212> DNA
<213> Meloidogyne incognita

<400> 103
ttggaccgtt aggattgtcg ccaaagaaaa ttggagaaga cattgcaaag gcaacacaag 60
actggaaagg cttaaagggt acttgcaaat tgactatcca aaaccgaatt gcca 114

<210> 104
<211> 255
<212> DNA
<213> Meloidogyne incognita

AKK110P1

<400> 104
 ccgcttctcg aattgtgaag gaattgaagg aacctcaccg agaccgcaaa aaagtcaaac 60
 acgtaaaaa cagtggaaat ttgacgatcg agcaaatat caacattgca cggcaaatgc 120
 gacctcgttc aatggcgaaa aaaattggaa gggactgtta aggaaattct tggcactgca 180
 caatctgttg ggtgtactgt tgatggacaa catccacatg atattgttga tgcaatccga 240
 agtgggaaaa ttgaa 255

<210> 105
 <211> 571
 <212> DNA
 <213> Meloidogyne incognita

<400> 105
 tttttttttt tttttttttt tgtcaacaat aaattttactc agaaaaatca tttacaatt 60
 taacacacat ttttaattcc ttaatactcc aaaaaacttc tcttctttat tccctcttat 120
 tctcccaatt catttaaagt ttcagttttg tgcggcgcca atgacgacgt tttgcattat 180
 agcgtatacg actgccagtt ttcattcgaa cccattgagg cagcgggtcga tttgttttag 240
 cagccttagc cagcttgccg ttgataataa acgttttggg tgcagccatt aaattgttga 300
 ctttatccaa aattgttttt ttgaaggcaa taaacaaatt taatttttct gctcaacaag 360
 tccatagcag ctcatctggt caacaatctc cctcatgccc ctcagtctcc agcgcttcct 420
 cttatgaatg tcaaaaacag cagcaacaac ccccagcaga accttgtgga ctttctttgg 480
 aagttcatca atctggtcat tcaacaacaa cccttccatc tccatgttct ttattacccc 540
 ctccctcttc tttacatcct ataatcatc g 571

<210> 106
 <211> 235
 <212> DNA
 <213> Meloidogyne incognita

<400> 106
 tgcttttatt tcaattcttc aaccaaaaaat taaatcttcc cttattttta ttacaattcc 60
 aatttttagc gcattagccc caactacttt agctgctaataaaaattgttt atgaggatgg 120
 agatagtgat ggacttgata tggctaaaag tatttttaaat tgaataaagg aaaaagaagc 180
 attttaaaaga aaatttagatg gaaatgctga agaaagaaaa aaattattta ttttt 235

<210> 107
 <211> 702
 <212> DNA
 <213> Meloidogyne incognita

<400> 107
 ttttttcaaa aaataattcg aattttgttc ttttttattt tgctacaaat aaaattttaaa 60
 tttgaaaaaa aaaaaaaaaa aaaaaaaaaa tgcgagaaga atccttgccg aaattgacgg 120
 ctctcaaatg gaggagtatc aacgtttctt cgatatgttt gaccgtggaa agaattggcta 180
 tattatggct actcaaatg gggtaattat gaatgctatg gaacaagatt ttgatgaaaa 240
 aactcttcgg aaattaatcc gaaaattcga cgcagacggc agcggcaaaa tcgaattcga 300
 cgaattctgc gctttggtat acactgtggc gaatactgta gacaaggaca ctttgccgaa 360
 agaattgaga gaagcttttc gtctctttga caaagagggc aatggttaca tctctcgtcc 420
 aacactcaaa ggattacttc acgaaatcgc cccagacctc agcgataaag acttggatgc 480
 cgcagtagac gagatcgacg aagacggaag cggaaaaatt gaatttgaag aattttggga 540
 gttaatggct ggagagactg attgaaattt taattagaat gactagaaaa ttgactaaaa 600
 tattttgcca ttaaattttg gaaagtgcga aaaattgcct ttctgagaat ttttattttt 660
 aacgtctaaa taatgaataa aatggatata aaaaaaaaaa aa 702

<210> 108
 <211> 423
 <212> DNA
 <213> Meloidogyne incognita

<400> 108
 aaaatttaaa taaaagacaa acaataaaat ataaattaaa taaataatat ttaaataaac 60
 acacaaataa actctccaaa cataattttt ttaaatttta ataacatttt gtcccatattg 120
 agaaagaaaa tgccaaagga gatgaagaac ttgttgaaga aaaaagttca aaaatatcaa 180
 ctctccatt tgctcgtcaca ttttctttca ttattccatt tgttgtaagc tcagtaactg 240

AKK110P1

```

cccccaattgt tgttgtagtc catggagaga aagcactttc cccattcgaa aatggtgaac 300
caaattgggtc aaattgttgc tgttggtgac ctcgaagtgc gttagaaaca gaacgaaata 360
aattatgagg ttgttggtgt tcctgacggt tttgattgtc tggagctggg tgaggatcac 420
caa                                                    423

```

<210> 109
 <211> 994
 <212> DNA
 <213> Meloidogyne incognita

```

<400> 109
ttttatTTTT tatttgaaaa taatcatcac attataatta atgggaaaaa gacaaaaaat 60
tagaacaggt gctggcgatc ttgtcacaac ccctggacct cttcataaac aaattgaaag 120
gtcaaaaacta gccaaagccga aattcaagcc tttaaaacgt tcaagagaag agcaaaaaaga 180
tgaaattgaa cttgtcgatc catcgttaaa gggcaaaatt attattaaag caaacaaaaa 240
attggaaaaa gatgttggtg tcaatgagga tggagaatct gataattctg aagaaattga 300
agaagaagaa gaagacggca atgaaaagtt ggatgttgat caattagtat caaaacattt 360
ggaagattta gatgaactaa aattggatga tggcgttgaa aatgtgCGaa agataataac 420
gaaattcaga taaaaataac aaagaaagtg ttataataaa agctgagttt gccgatatcg 480
acccaaaaat tgttgatctt ttacagaaa ttggtcaagt tttaaagaaa tatagaagtg 540
gacgtattcc caaagctttt aaagttattc caactttggt tgattgggag aaaattatcg 600
aattaactcg cccagatgat tggtcggcag ctgcaatggt acatgctacc aaaatatttg 660
cttcaactgc tacccctact caatgccaaa gggtttataa tttgattttg ttgccacgta 720
ttcgagatga tattgacgga ttaaaaaatt acatttccat atgtatcaat gcttatttaa 780
agcattgttc aaaccagctg catttttcaa aggaatcctt ttgccgcttt gcaaatcgaa 840
caatttttct cttcgagaag ctgttggtct tgcttctatg cttcgtaaag cctccatccc 900
tcaattacac gcggcccgag cattgttgag tatttcttgt ttagaatata cttcttcaag 960
ggcttatatc cttcaagcat tgatagaaaa gaat                                                    994

```

<210> 110
 <211> 476
 <212> DNA
 <213> Meloidogyne incognita

```

<400> 110
tttaaacact taaaaatacc ttcaaattta ttttagaacc tttttgccat taaaaaaaat 60
tttattttcga aaaaatggct gagaatatag aagaaatcct tgccgaaatt gacggctctc 120
aaattgagga gtatcaacgt ttcttcgata tgtttgaccg tggaaagaat ggctatatata 180
tgggccactca aattggggta attatgaatg ctatggaaca agattttgat gaaaaaactc 240
ttcgaaaaatt aatccgaaaa ttcgacgcag acggcagcgg caaaatcgaa ttcgacgaat 300
tctgCGcctt ggtatacact gtggcgaaata ctgtagataa ggacactttg cggaaagaat 360
tgagagaagc ttttcgtctc ttcgacaagg agggtaatgg ttacatctct cgtccaacac 420
tcaaaggatt actccacgaa atcgccccag acctcagcga taaagacttg gatgcc 476

```

<210> 111
 <211> 189
 <212> DNA
 <213> Meloidogyne incognita

```

<400> 111
cgaagacgga agcggaaaaa ttgaatttga agaatttttg gaattaatgg ctggagagac 60
tgattgaaat ttaatttaga gatgaataaa aaattaacta aaatattttg ccataaaaatt 120
ttggaaaagt ccaaaaattg cctttttgag aattttttatt tttaacgtct aaataatgaa 180
taaattggat                                                    189

```

<210> 112
 <211> 164
 <212> DNA
 <213> Meloidogyne incognita

```

<400> 112
ttgaggaaat ttaatttttt aaacaaatat aataattacc aaacaacaaa aaagaatccc 60
aaaaacaaca tttttaaatc aaatgacaga catatatatt caataacgat gtgtggattt 120
tctttttttt taaataatta acatctttaag cctgctattt cttc                                                    164

```

AKK110P1

<210> 113
 <211> 539
 <212> DNA
 <213> Meloidogyne incognita

<400> 113
 cagctttctg cgcagatttg gtaacctttc caccagcttc gaccttctcg acggccttga 60
 taacaccaac agccacagtt tgacgcatgt cacgaacggc gaagcgtcca agaggagcgt 120
 agtcagtaaa agcctcaaca cacattggct tggittggaat taagtcgaca ataccagcat 180
 ctccagtcctt caaagccttt ggattgtctt caaccttctt tccagttcga cggtcgacct 240
 tctctttaag ctacgcgaac ttgcaagcaa tgtgagcagt gtgacagtca agaacaggcg 300
 tgtagccagc agcaatctgc ccaggatggg tcatgatgat aacctgagca gtgaattgct 360
 tggctctcct tgctgggtca ttcataagat cagaagtgc tgaaccacgt cggatgtcct 420
 tgacagagat gttcttaacg ttaaatccaa cattgtcttc aggaacagct tcaggggagag 480
 actcgtgggtg catctcaaca gatttaactt cagtagaaat tccttcagga gcaaaggta 539

<210> 114
 <211> 314
 <212> DNA
 <213> Meloidogyne incognita

<400> 114
 gtttttaatt ttagaaaatg tctacagaâa cagaaaagga tttagaacgt tgggaggatg 60
 tccgtcgatt tactgagatt ggttcttcta aatttgccca tcccgccttt gttccaagcc 120
 cggagaatct tgaaagagta aggaaatgtc cagttttggg tggttggtgct ggtgggcttg 180
 gatgtgaaat ttgaaaaat ttggccttat caggatttca aaatattgaa gttattgata 240
 tggacacaat tgacctttca aatctcaaca gacagttttt gtttcgtgaa cacgatgttg 300
 gcttatacaa agca 314

<210> 115
 <211> 200
 <212> DNA
 <213> Meloidogyne incognita

<400> 115
 ttcgaagacg tggttaaagga tgcgtcttta ctgcacataa ttgtaaaata caagataaag 60
 gacttgactt ttatgggcaa ttttcaatta taatttggg actagattct attgatgctc 120
 gaagatgggt aaacgccaca gtgtgttctt tggtcgaatt tgacgaagaa aacaagccac 180
 ggccaggcac aattattcca 200

<210> 116
 <211> 471
 <212> DNA
 <213> Meloidogyne incognita

<400> 116
 tttggtcgaa aaaagactgc tactgctgtg gcatattcca aaaagggaaa aggattaatc 60
 aagggcaatg gccgtccttt agaatttttg caacctgaaa ttcttcgtat taagctacaa 120
 gagccattgt tgattgtagg aaaggacaaa tttgctggaa tggatattcg catccgtgtc 180
 aaagggtggtg gtcattgtgc acaaatttat gcaattcgac agtcaattgc taaagttttg 240
 gtggcctatt accagaaaaa cgtggatgag caaagcaaga aagaattgaa ggatcaactt 300
 gttgcttatg atcgtaattt gcttggtggc gatccgagac gtcacgagcc aaagaagttt 360
 ggaggacctg gtgctcgtgc tcgttatcag aaatcttatc gttaagaagt atgaaattat 420
 aaaattgtgt gttacgaatt aattgttatt ttgttgggat aaatntgaat a 471

<210> 117
 <211> 593
 <212> DNA
 <213> Meloidogyne incognita

<400> 117
 gaattcaaaa aatattaaaa ttgtttaata taatttctaa aatgaagcca aaggttggaa 60

AKK110P1

ttaacggatt	tggacgtatt	ggacgtcttg	ccctgcgtgc	agcgggtcgag	aaggatactg	120
tccaagtgt	ggctgtcaat	gacccgttca	ttgatcttga	ctatatgggc	tatatgttta	180
actatgattc	cacccacgga	cgcttttaag	gaaagattca	agcaagcaat	ggaaaatttg	240
tagttgagaa	ggagggaaaag	tctactcata	ctatcaaagt	tttcaacttc	aaagaacctg	300
aaaagattga	ctgggcaggt	tctggtgctg	atcttgttat	tgagtcgact	ggagttttta	360
ctactaccga	gaaagcttct	gctcacttga	agggcggagc	caagaaagtg	gttatctccg	420
ctccatctgc	tgatgctcca	atgtttgtgg	ttgggtgtta	tgaggacaaa	tatgatcctt	480
ccaagcatca	tatcattagt	aatgcttcct	gcactactaa	ttgtcttgct	cctcttgcca	540
aggttataaa	tgacgagttt	ggcataattg	aaagttgaat	gactactgga	cac	593

<210> 118

<211> 576

<212> DNA

<213> Meloidogyne incognita

<400> 118

gaattccgag	tttttttttt	ttttttttta	aacaaaaatt	aaaagattta	tcgccatcct	60
ttgcccagca	tttgcctgcc	atctttttgt	gcacaataaa	tttttttgta	atctttgggg	120
tgagggggaa	gtaaaatgaa	agaagggaga	gagatatgaa	ttggagggtt	ttttgttaaa	180
ataaattttt	ttttcttgaa	aattcttccc	gtttctgagc	tttttcgtct	tttttcaatt	240
ttcgtttgct	gaaataactaa	actttacaat	ttggttaggt	tctattttgtg	aaacataaat	300
atctccatta	tcgctgattg	caagggcatg	ggcgttttcg	agaccctttg	caaagctatt	360
agcccttccct	gtgttcataat	ccattacgaa	aacttgggat	tctaattgac	tgcccttgatc	420
ttgattgggtg	acgccgacga	ggaagtgttc	tttctctcgg	atagcaaaga	ctcgcccaat	480
atttttcagcc	tttgtgaaga	aagtgcctgt	ggggacgtaa	gcacgtctat	gttggtgttg	540
agcgcttctt	aatccagcag	aaaagcattg	aatacgc			576

<210> 119

<211> 559

<212> DNA

<213> Meloidogyne incognita

<400> 119

acgcagagta	agttgagatc	ttcaataagg	gttagagagt	gtggtacgag	gaattctcca	60
tttttggttg	tttctactgga	gtcaggcttc	ccaaattgac	tgagcaattt	cccctccttg	120
tcaaacttca	ttattcggct	attacagtta	ccatctgcca	cgaaaaactc	tcctgtactg	180
gcaatagcaa	cgctctgtag	tttgcaaaaa	tgtttgctat	ctgtccctgg	aacaagcttt	240
tcgcccacaac	tcataattaa	tttaaaatcc	ttgtcaagtt	tggtggacttg	atgacttcca	300
acgtcagtaa	cccaactatt	gccgtgggca	tcgattgtta	gtccatgagg	catgtaaaac	360
atgctttttc	cgtattcttc	caagactgcc	cctgattccg	tgtctataac	agcaattggt	420
gtggttgaaa	tgatgcccg	ggatctgttt	aggtggttgt	tctcatcaaa	cgaaaattca	480
tcccacaactc	tgtcagatcg	gtgaaaaaga	acaagtcgat	tcaatggatc	caatgcaata	540
cccggagctt	gcccataat					559

<210> 120

<211> 366

<212> DNA

<213> Meloidogyne incognita

<400> 120

tttaagaatt	ttttaaaaaat	taaaacttgg	actagatttt	aataaaaatgt	cagctccacg	60
tagtggtgct	agcgggtgtg	gtgctgctgt	tatgaataag	caagcaagta	aatacaatga	120
agttgaagga	gaactccttc	tttaattggat	taagaaagtg	acaggcgaaa	atattgctat	180
aaacgggaact	agggaaaatt	ttgtgaaaca	attgaaagat	ggaaactctgc	tctgcaaat	240
tgctaacaaa	attgtgccaa	attcaatcac	aaaggcacag	gcaaaaccga	acagcacatt	300
ccaatatatg	agcaatttgg	agctgttctt	aacattttatt	tcaagccaag	gagtccttag	360
ggagga						366

<210> 121

<211> 661

<212> DNA

<213> Meloidogyne incognita

<400> 121

AKK110P1

ttagttgaat	ctcgtgacct	ctactctgtt	tgtatgacat	taaattctct	tgcccgcat	60
ttggaacgtc	aaggaaaaac	tcattccagag	cagggttaagt	cgtcagaaat	tcttaatttg	120
ggtagctggag	accaagtgcg	ccttcgtgtt	taaagatggg	aaattgaaag	aattttgggt	180
aaacataata	aaaagacatt	ttatggcaat	aaaaaaatgt	caaaaaagct	tgtcttttaa	240
atattttggc	aaaacatttt	actttcacaa	aatttttaaaa	taaatttatg	aagattgttc	300
cgctactttc	atcattttccg	atcgaccttt	gttgttttct	aagttcggtg	gccaaagaaa	360
ggatatgtaa	aattgaatta	tgaataaaaa	taaatcactc	aatcagaggg	attgttagtc	420
tctcactttc	tcctctttac	ccattggcta	accagcttta	aggatttttt	ccataagttc	480
aagggtgtacg	taaatcgaat	accgactgtg	gtatcttaat	ttttccatga	aattctccaa	540
taaaaaaaaa	ttttttttat	tttttttcca	taatgctatc	tatatatttt	gcttttaatc	600
ttttttggct	atcaggcctt	aaaatagtaa	atatacttat	attaatat	tatttccttt	660
a						661

<210> 122
 <211> 173
 <212> DNA
 <213> Meloidogyne incognita

<400> 122						
ggagagtttt	tcgtggcaga	tggttactgt	aatagtcgaa	taatgaagtt	tgacaaggat	60
gggaaattgc	tcagtcaatt	tgggaagcct	gactccagtg	aaacacccaa	aaatggagaa	120
ttccttgtag	cacactctct	aaccctcatt	gaagatctca	acttactttg	tgt	173

<210> 123
 <211> 584
 <212> DNA
 <213> Meloidogyne incognita

<400> 123						
cgcattcaat	gctttttctgc	tggattagaa	ggcgctcaac	accaacatag	acgtgcttac	60
gtccccacag	gcacttttctt	cacaaaggct	gaaatatattg	ggcgagtctt	tgctatccga	120
gagaaagaac	acttcctcgt	cggcgtcacc	aatcaagatc	agggcagtc	attagaatcc	180
caagttttcg	taatggatat	gaacacagga	agggctaata	gctttgctaa	gggtctagaa	240
aacgcccattg	cccttgcaat	cagcgataat	ggagatatatt	atgtttcaca	aatagaaccc	300
aaccaaattg	taaaatttag	tatttcgaca	aacgaaaatt	gagaaaaaaa	aaaaaaaagc	360
tcagaaacgg	gaagaatttt	caagaaaaaa	tttttttacc	aaacaaaaaa	cctccaattc	420
atatctctcc	cttctttcat	ttttccttcc	ccttctcccc	aaaaattaca	aaaaatttta	480
ttgtgcacaa	aaaaatgggc	gggcgggcga	atggctgggc	aaaggatggc	gataaatctt	540
ttaatttttg	aaaaaaaaaa	aaagaattcg	aattatatgg	ccta		584

<210> 124
 <211> 650
 <212> DNA
 <213> Meloidogyne incognita

<400> 124						
gtttaagaca	attaaaacgt	ttattttcta	caatcaaaac	aaatatggct	gttctctccg	60
atgttatcga	gaagatcgag	gctgggtaca	aaaagttgca	ggaggcaccg	gagtgcaggt	120
ctcttctcaa	gaagtacttc	acgaagggaag	ttatggacca	gtgtaaagg	ctcaaaacta	180
agcttggtgc	gaacttgctt	gatgtgatcc	actctggagt	tgcgaatctc	gatagcgggtg	240
ttgggtgttt	tgcgcctgat	gctgagcttt	acactctctt	caaaccgctt	tttgacccga	300
ttattcagga	ttaccacaat	ggatttggac	ctgaccagaa	gcagccgcaa	actgacttgg	360
gtgagggaaa	gactcagctt	ttgcctgatc	tggatcctga	gggtaaaattc	atcaactcga	420
ctcgtgttcg	atgtgggcgt	tctcttcagg	gatatccgtt	caatccgtgc	ttgactaaag	480
agaattatac	ggaaatgcat	gacaaagtta	aaggggtttt	tgagcagctt	aagtctgatg	540
ctgagcttgg	tggcacctat	tatcctttgg	aggggaatgac	caaagaggtt	caaactcaat	600
tgatcaagga	tcacttcctc	ttcaagaag	gagaccgctt	tttgcaagct		650

<210> 125
 <211> 1013
 <212> DNA
 <213> Meloidogyne incognita

<400> 125

AKK110P1

tttttttttt	tttgatgttt	ctaatttttt	tgggcaatat	ttaatattat	ttttaattat	60
ttaaattttt	tctttatttt	ttaaaaaatt	atttctttaa	tttattcttc	tcctcttcgt	120
gttttgaatc	aaataattaa	attttaaatt	atttaaacag	ctacacgagg	cctcagcctc	180
ccccgttgca	ttcaaattgg	tcggcacggt	tggcgatgat	aattttat	tttaggtaat	240
tttggtgaga	aaatattttt	aaaggtaata	atgtcctttt	ggacaattaa	aaaaaaactc	300
gaggagagag	tgaatatttt	tacaaattat	ttgaagagca	gccagcctat	tgttatcaac	360
aaaaaacctt	caaaatgcca	gaaaatgatt	atgatgagga	ggaggcgcca	aacgccacga	420
tggaaacaaa	ggtagcttca	ggtggacagc	caaaacgctg	ttggaaaatg	gacattatcc	480
cagctgcgcc	agactgatgg	tataattcca	tcccaggccg	gttggaaaca	gggagactcc	540
caaaagtiga	tgaccaattt	tggtactcca	cgtaacacaa	caaccaaaat	tcgtgctgaa	600
tgcttgctg	atgtgcctga	agaaattgct	cttaaaagtc	acggtgaagt	acgcctccaa	660
tccggtacta	accgttttgc	ttcgcagaag	ggaatgggtg	gatttggtac	tggaactgac	720
ttatgcagag	aaggagtgtt	tgtgagtcaa	gacccagccg	atttatagcc	cctcccagaa	780
gagataatcc	gtgctagcga	tggaaattgt	cgtctccaat	ccggtacca	caaatctgac	840
tcccaaaagg	gaatggtcag	cttcggtaca	aaccgacgag	aaactacaag	aatgaaagac	900
accaaacatc	cggaatacaa	ccacgaagtt	aacattgacc	aaagcgaaat	tcctttgcaa	960
tctggtacaa	acaaattcgc	atcccaaaag	ggaatgacca	gcttcggtac	aaa	1013

<210> 126
 <211> 80
 <212> DNA
 <213> Meloidogyne incognita

<400> 126
 tggtggacac tgctcaccca gaatacagtc acgaaagcag catcgatcaa acgagcattc 60
 cttaccaaat gggatcaaat 80

<210> 127
 <211> 585
 <212> DNA
 <213> Meloidogyne incognita

<400> 127
 agggaaatgac ttgcttttga cagccacggt gggagggtgct tgacccgagc attagctacc 60
 agaaccgtaa atcacaaagga atggtccgct tccaatccgg aacaaaccgg gtcgcctcgc 120
 aagcgggcat gacagggtttt ggaactccaa ggaacacaaac atacgaggcg gactctggcg 180
 aacttccata cgaagatatg aagaagtcag aaacgataat tccatcccag gccggttga 240
 ataagggaga ctctcaaaag ttgatgactg gatttggtac tcctcgtgac gttaaaggca 300
 aacatttgaa gcgtatttgg gagttggaat acccagagga ggctgaaatt tcgttggaac 360
 gacttttaag gaatttttaga agagaagaaa gaaaagagaa atttagtgga aggaaggcaa 420
 cgacatttga ctctacaatt gacacacacc ttttcacaca tttaaaaaat acattaaaaa 480
 aaaatttttt ttggcttttt ggcttgctcc tatttttttc ccccatcatt ctccctattc 540
 tctcatttgg atgcaaactg gaatttttaa aaaaaaaaaa aaaaa 585

<210> 128
 <211> 287
 <212> DNA
 <213> Meloidogyne incognita

<400> 128
 catctggaga aacgttgagg caatacatcg ttattggccg taaacttcct acagagaatg 60
 agccaaatcc aaaactttac aaaatgcaaa tttttgccag taatcatgtt gttgctaaat 120
 cgcgtttctg gtactttact agtatgttgc gtcgtgttaa gaagactaac ggagagattg 180
 tttcgtgtca ggagggtttt gaaaagaaga taggctctgt aaagaattat ggaatttggc 240
 ttcgttatga ctctcgaacc ggtcatcaca acatgtaccg tgaatac 287

<210> 129
 <211> 175
 <212> DNA
 <213> Meloidogyne incognita

<400> 129
 gctgtcactc aggcattatcg cgacatgggt gctcgtcatc gtgctcaagc cgatcgaatc 60
 caaataatca aggttcaacc gatcaaggct gccgattgca aacgtactgg agttaaacag 120

AKK110P1

ttccacaact cttcaatcaa gtttcctttg ccgcatcgtg tgaatgacaa acgtc 175

<210> 130

<211> 599

<212> DNA

<213> Meloidogyne incognita

<400> 130

acttttgttt ataatcacat ttgcattact ttccgtccat ctttctttga gacagaattt 60
 aaagggttcac cttctaagta aggattgtag cggctgtatg attgatgttg cttttgttgg 120
 ggagcaatag aacgcttgcg tcgccgaggc tcctcagccc tagtaacgtg aaatttcttt 180
 gcaatcatcg atttgtgtag tccatttttg gctaagacct gttctaagtc ttgttcatat 240
 tgttcagaat tgctttttga ttgacagtta aacatgtgtt cttggtcaca aaggcattgc 300
 tgattggcct ggtagctacg cgagaaatcg gcggtgttat caaactcctc caaacatcca 360
 tctcgactgg agtatccac agggcaggga tttggagggt cacaatatgc tggcaaaaca 420
 ttgtcactct taatctcttg gcggtgtgaa aattcagatt ctggatggag ttgttggtct 480
 ctttcaccgg cacctcctgt cataaattta tgtccaaacg caatgggccc ggaagcactt 540
 tcaatgtcac gagaaatcaa gtcgattaat tgtgaatgag gaaatatagg ctccccaga 599

<210> 131

<211> 466

<212> DNA

<213> Meloidogyne incognita

<400> 131

gaagattgga tttattggcg ctggaaagat ggcacaggca ttggccagag gactaataaa 60
 ttctggacgt tatccttcac aaaatttgat ggctagtgtc cctaagactg atgtctcttt 120
 attggaggat tgcaagaggc ttgggagtaa tacagcacat gataatgcac aagttgctcg 180
 tgaaaatgat gtggtgatta tagcaggtaa accaactatt gtgtctaaag ttgcttcgga 240
 aattgcacca gccatccgcc gagatcatgt acttatttct atagcattgg gcatcaccat 300
 acgctacatt gagcagtaat tgacttcaga atcccgaatt gttcgtgtaa tgccagatac 360
 tcctgtagggt ggtaggagca ggctgctgca gccatatatc attgggatca gcattgtcag 420
 gatagggtgat gcccagatag ttcaagatct tctgataacg ctggggg 466

<210> 132

<211> 266

<212> DNA

<213> Meloidogyne incognita

<400> 132

atgaaattcg agttctttgc atcaaggccc gtgaaatttt tctttcgcga cctatttttgc 60
 tggaaattgga agcgccgttg aagatttgtg gcgatattca cggatcaatac aacgaccttt 120
 tgcggctttt tgaatatgga ggttttccgc ctgaagcgaa ttattttatt ttgggtgatt 180
 atgtggatag aggaaagcag agcttggaga cgatttgttt gctgttggcc tacaagatca 240
 aatcccccca aaattctttt tgctga 266

<210> 133

<211> 308

<212> DNA

<213> Meloidogyne incognita

<400> 133

tctatcaacc gaatatatgg attttacgat gaatgcaaac gcagattttc tataaaattg 60
 tggaaaacat ttactgattg cttcaattgt ctgccaatg ctgctgtgat cgatgagaaa 120
 atattttgtt gccatggagg tttgtcacca gatttgcaga atatggagca aattcgaaga 180
 attatgcgac cgacggatgt gccagatata ggtcttctct gcgaccttct atggtctgat 240
 ccagaccaag atgtccaagg attgggagaa aatgatcgtg gggctctttt cacttttggg 300
 ccagatgt 308

<210> 134

<211> 335

<212> DNA

<213> Meloidogyne incognita

AKK110P1

<400> 134
 taaatttagt ttcttttctt ccatctcttt ttatgttttg aaagagtgtg ccaaaaacaaa 60
 tggccgcccg tgatggaaga agcaggcaaa attatttaca agaacattca attcctcaac 120
 tttttgaggg tttaatgact ggacttatat acaatcaacc aatcgatcct attcaatttt 180
 tggagaatgc aatagctaaa cttcgaaaaa atcctgatct tccattaaag tgggatactt 240
 ttataagtgt ttcgccctcaa caacagcaac aacaacagac gagaatgaat actggagaaa 300
 atgcagtttc ttataaacia agcactccta tcgaa 335

<210> 135
 <211> 506
 <212> DNA
 <213> Meloidogyne incognita

<400> 135
 tttttttttt tttaaaaatc aacagattta ttcaagtgcc tcgggcaaatt aacaacaaac 60
 atccacaaac ataattattat tgaacttttc ctttttaaaa cttatcaaag gccttctttg 120
 ttcttgagac ttgatcacc ttcaaaacat taaaacgaac agttttactc aaaggcctgc 180
 attcacccgat cgtgacaata tcaccaatag agatatcacg gaaacatggc gaacagtga 240
 cggacatgtt tttgtgacgt ttctcgtatc gacgatattt cggaacaaag tgcaataat 300
 cacgccgaat gacaattgtg cgctgcattt tgttcttgat aacaacacca gtcaaaatac 360
 ggccacgaat tgaaacattt ccagtgaag gacacttttt gtcaatataa ttgccttcga 420
 tagcctcgcg tggagtttta aatcctaacc caacattctt ccaataacga tccttatttt 480
 tcggctnttt gccaatccct tgcgtc 506

<210> 136
 <211> 230
 <212> DNA
 <213> Meloidogyne incognita

<400> 136
 aattcctcaa actctgccct ggctgtcctt ctcaaaacga caccctcgct ttattatcac 60
 ctccagtcac ctacgaaaat tctttgcgag atcaaggag taattcgaca ttatggattc 120
 ttttgttggt ttttaattgt ttatttttgc tactaatttt ccttctaatt gccgcctacc 180
 tccgttgtcg cttttttggc tccgccccct acaaaaacca gttccgtcgt 230

<210> 137
 <211> 216
 <212> DNA
 <213> Meloidogyne incognita

<400> 137
 acaaatcac aacaacaaaa tcattgttta ccccaaatat ccaaaagttc tcctccaact 60
 tcaactcgctg tttgtaccaa ctctactagc tgtaattcgt ccttagctgt gccgttaatt 120
 tctagtgaat cggaagaaag tgatgaacaa caaagacgg gggaaatggac aaatctaaca 180
 ttattaatta tttattctca tgattgtaaa ttgcat 216

<210> 138
 <211> 395
 <212> DNA
 <213> Meloidogyne incognita

<400> 138
 atgcattcct gaagcaattt tgggtatgga cattgtatgc caagcaaagt ctggtatggg 60
 gaagacagct gtatttgtgt tggcaacact ccaacaattg actccagttg acgggacggg 120
 ctctgttctc gttatgtgtc acactcgca acttgctttt caaatttcaa aggaatatga 180
 aagatttagc aaatatatgc ccggaactaa ggtttcgggt ttctttgggt gtatgccgat 240
 caagaaggac gaggagactt tggctaagaa cactccgcac attgttgttg gcactccagg 300
 gcgtctgctg gcgttgggac gtacaggaca attgaagctg aaaaacatca aattcttcgt 360
 tttagacgaa tgtgacaaaa tgattgggga cgctg 395

<210> 139
 <211> 591

AKK110P1

<212> DNA

<213> Meloidogyne incognita

<400> 139

gaattcggcg	ttgtctcggt	gtccacgctc	aatttcaccg	aaatTTTTtg	ggcaggcgctc	60
ctccacacca	aactctgggt	cattgacaac	cggcactttt	atctgggttc	agcaaacatg	120
gactggcagt	cacttactga	agtcaaggaa	atgggtctta	tgctgttgaa	ctgctcctgt	180
ttggcgtggg	aactgagcaa	aatatttgcg	atttactggc	ggattggaca	gaatcacaat	240
cgcttgcccc	ctgtttggcc	agttttattta	caatcaaaaat	tcaacgctca	acacccaatg	300
gaaattcatt	ttggacctga	gccctcgcac	acgtacattt	cgactcgcc	tgagaagttg	360
aacccaaagg	gcagagaaca	cgacctttcg	gccatatgct	catgcatggg	aaaagccaac	420
gaatttggtc	gaattgcggt	aatggattat	attcctgcaa	caatttacat	gccgaatggt	480
aacaacatat	attggccatc	gatcgatgac	gcgataagaa	cggcagctta	tcgggggtgtg	540
aaagttgacc	tttgggtgagt	ctgtggcccc	atttgaatga	acgagcgatt	t	591